

Ministry of Regional Affairs and Agriculture, Republic of Estonia

**Regulation (EU) 2017/1004 of the European Parliament and of the
Council of 17 May 2017**

on the establishment of a Union framework for the collection, management and use of data in
the fisheries sector and support for scientific advice regarding the common fisheries policy
and repealing Council Regulation (EC) No 199/2008 (recast)

Commission Delegated Decision (EU) 2021/1167 of 27 April 2021

establishing the multiannual Union programme for the collection and management of
biological, environmental, technical and socioeconomic data in the fisheries and aquaculture
sectors from 2022

**Commission Implementing Decision (EU) 2021/1168 of 27 April
2021**

establishing the list of mandatory research surveys at sea and thresholds as part of the
multiannual Union programme for the collection and management of data in the fisheries and
aquaculture sectors from 2022

**Commission Implementing Decision (EU) 2022/39 of 12 January
2022**

laying down rules on the format and timetables for the submission of national work plans and
annual reports for data collection in the fisheries and aquaculture sectors, and repealing
Implementing Decisions (EU) 2016/1701 and (EU) 2018/1283

**ESTONIAN Annual Report on data
collection in the fisheries and aquaculture
sectors**

2023

Version 3

Tallinn, 30.05.2024

CONTENTS

SECTION 1: GENERAL INFORMATION

5

DATA COLLECTION FRAMEWORK AT NATIONAL LEVEL

5

TEXT BOX 1A: TEST STUDIES DESCRIPTION

6

TEXT BOX 1B: OTHER DATA COLLECTION ACTIVITIES

7

SECTION 2: BIOLOGICAL DATA

8

TEXT BOX 2.1: LIST OF REQUIRED SPECIES/STOCKS

8

TEXT BOX 2.2: PLANNING OF SAMPLING FOR BIOLOGICAL VARIABLES

8

TEXT BOX 2.3: DIADROMOUS SPECIES DATA COLLECTION IN FRESHWATER

11

TEXT BOX 2.4: RECREATIONAL FISHERIES

13

TEXT BOX 2.5: SAMPLING PLAN DESCRIPTION FOR BIOLOGICAL DATA

14

TEXT BOX 2.6: RESEARCH SURVEYS AT SEA

16

SECTION 3: FISHING ACTIVITY DATA

26

TEXT BOX 3.2: FISHING ACTIVITY VARIABLES DATA COLLECTION STRATEGY (FOR INLAND EEL COMMERCIAL FISHERIES)

26

SECTION 4: IMPACT OF FISHERIES ON MARINE BIOLOGICAL RESOURCES

27

TEXT BOX 4.2: INCIDENTAL CATCHES OF SENSITIVE SPECIES

27

TEXT BOX 4.3: FISHERIES IMPACT ON MARINE HABITATS

31

SECTION 5: ECONOMIC AND SOCIAL DATA IN FISHERIES

33

TEXT BOX 5.2: ECONOMIC AND SOCIAL VARIABLES FOR FISHERIES DATA COLLECTION

33

SECTION 6: ECONOMIC AND SOCIAL DATA IN AQUACULTURE

34

TEXT BOX 6.1: ECONOMIC AND SOCIAL VARIABLES FOR AQUACULTURE DATA COLLECTION

34

SECTION 7: ECONOMIC AND SOCIAL DATA IN FISH PROCESSING

34

TEXT BOX 7.1: ECONOMIC AND SOCIAL VARIABLES FOR FISH PROCESSING DATA COLLECTION

34

ANNEX 1.1 - QUALITY REPORT FOR BIOLOGICAL DATA SAMPLING SCHEME

36

ONSHORECOMMERCIALCOASTAL

34

SELFONSHORECOMMERCIALCOASTALSALMON

38

ONSHORECOMMERCIALPELAGIC

40

ONSHORECOMMERCIALPELAGICGOR

43

ATSEACOMMERCIALNEAFC

45

ATSEACOMMERCIALNAFO

47

PARRELECTROFISHING

49

SMOLTTRAPPING

52

ADULTCOUNTING

54

CFSEEL

56

FRESHWATER_EELFYKES_SCIENTIFIC

58

FRESHWATER_EELFYKES_COMMERCIAL	
.....	
60	
RECREATIONAL_CATCHREPORTS	
.....	
62	
RECREATIONAL_INTERVIEW	
.....	
64	
BITS_Q4	
.....	
66	
BIAS	
.....	
69	
GRAHS	
.....	
71	
SPRAS	
.....	
74	
EFLS.....	76
GORFS.....	78
CFS.....	80
ANNEX 1.2 - QUALITY REPORT FOR SOCIOECONOMIC DATA SAMPLING SCHEME.....	83
SOCIOECONOMIC FISHERIES DATA: CENSUS.....	83
SOCIOECONOMIC FISHERIES DATA: PSS.....	85

SECTION 1: GENERAL INFORMATION

Data collection framework at national level

General comment: Use this text box to describe how data collection is organised in your Member State (institutions involved, contact information) and in which regional coordination groups (RCG) your Member State participates.

The Ministry of Regional Affairs and Agriculture (MoRAA) is responsible for the implementation of the data collection in the fisheries and aquaculture sectors and coordinates the European Commission objectives between the data collection authorities.

Suur-Ameerika 1, 10122 Tallinn, Estonia

Katariina Kurina (national correspondent): katariina.kurina@agri.ee

<https://www.agri.ee>

The Estonian Marine Institute of the University of Tartu (EMI-UT) is the main contractual partner of the Ministry of the Environment in the implementation of the data collection program and has been doing this work since the beginning of the commitment. Estonian Marine Institute is responsible for all scientific surveys under the data collection program and its researchers have the obligation to take part in relevant ICES working groups and other scientific working groups, which assess the state of fish stocks and marine environment, develop methodologies and provide assessments.

Mäealuse 14, 12618 Tallinn, Estonia

Redik Eschbaum (responsible person): redik.eschbaum@ut.ee

<https://mereinstituut.ut.ee/et>

Estonian University of Life Sciences (EULS) assess the state of the eel stock in inland waters and participate in the work of the ICES Eel Working Group (WGEEL).

Kreutzwaldi 1, Tartu 51006, Estonia

Priit Bernotas: pbernotas@emu.ee

<https://www.emu.ee/>

Statistics Estonia (SE) has been collecting economic and social data on the fishing fleet since 2019, before that it was the obligation of EMI-UT. SE is the main data competence centre in Estonia and is also responsible for coordinating the system of classifications and for data governance in Estonia.

Tatari 51, 10134 Tallinn, Estonia

E-mail: stat@stat.ee

<https://www.stat.ee/et>

The Agriculture and Food Board (AFB) compiles catch and effort data and first sale price data of commercial fishery.

Estonia participates in the following regional coordination groups: RCG Baltic, RCG North Atlantic, North Sea & Eastern Arctic, RCG Long Distance Fishery (since 2022) and RCG on Economics Issues.

Link to the national data collection website:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm>

(max. 1000 words)

Text Box 1a: Test studies description

General comment: This text box fulfils Chapter II, section 1.2 of the EU MAP Delegated Decision annex.

Name of the study: Pilot study on timing of GRAHS survey

1. Aim of the test study

Aim of this test study is to get more stable input data for Gulf of Riga herring population assessment.

For that timing of the survey needs to be determined and either the Gulf of Riga Acoustic Herring Survey (GRAHS) accordingly amended or Baltic International Acoustic Survey (BIAS) survey could be extended to Gulf of Riga.

2. Duration of the test study

2022-2027

3. Methodology and expected outcomes of the test study

IBAS methodology is used for both BIAS and GRAHS surveys, but there are differences in survey effort and timing. GRAHS has 2-3x times more trawling and acoustic tracks. On time wise BIAS is conducted in September/October, GRAHS in July/August. This test study aims to assess the suitability of either extending BIAS survey to cover also the Gulf of Riga or postponing the GRAHS survey so it takes place later at the year. It was recommended by ICES WGBIFS to try this approach to avoid fluctuations in current Gulf of Riga survey results. Expected outcome is more stable assessment of Gulf of Riga herring population with hopefully new time series of results. Currently it is presumed that survey time later at autumn could give more stable results compared to survey conducted in August, but it needs to be tested.

This test study is conducted together with Latvia

(max 900 words per study)

Name of the study: Pilot study on timing of GRAHS survey

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

Second year of the test study was affected by rough weather conditions, which limited suitable survey period of rented vessel. Even the time difference of 15 days, which is usually considered having negligible effects in the BIAS survey of Baltic Sea, resulted with considerably different results in Gulf of Riga. In 2024 it is decided to make the survey with minimum time difference in any event to test whether the differences in results are solely from time differences in previous surveys.

Achievement of the original expected outcomes of the study and justification if this was not the case.

The test study is planned to be conducted on several years before the results and decisions can be made.

Incorporation of study results into regular sampling by the Member State.

Currently it is too early to incorporate the study results into regular sampling scheme. This matter will be discussed after the finalization of the study.

(max. 900 words per study)

Text Box 1b: Other data collection activities

General comment: Use this text box to provide information on other data collection activities that relate to your EMFAF operational programme and need to be included in the work plan and the annual report. Describe activities that are funded by the DCF but fulfil objectives under other EMFAF priorities, like marine knowledge, or activities funded by the DCF, but without a direct link to the EU MAP specific requirements or WP template tables, like freshwater fisheries. You can also include one-off specific studies for a particular end-user need that do not enter the regular data collection.

Name of the study: RCG Secretariat

1. Aim of the data collection activity

Follow-up for the Project SecWeb (Mare 2020-08) to have a long-term supportive structure for RCGs on administrative side. Functioning secretariat that gives administrative support for RCG and ISSG chairs and manage the RCG web page (<https://www.fisheries-rcg.eu/>)

2. Duration of the data collection activity

Starting from 2023

3. Methodology and expected outcomes of the data collection activity

A detailed description of the secretariat functions, the implementation of the secretariat, the content of the website, the building blocks of the website and the business model for the provision of Secretariat role and website continuation (updating & maintenance) will be provided at the end of Project SecWeb in 2022.

(max 900 words per activity)

Name of the study: RCG Secretariat

Brief description of the results (including deviations from the plan and justifications as to why if this was the case). Achievement of the original expected outcomes of the study and justification if this was not the case. Incorporation of study results into regular sampling by the Member State.

During 2023 the RCGs Secretariat services and tools were financed by a short-term/low value contract by the individual Member States (23 out of 26 MS) together with a European Commission service contract running from the end of August 2023 to end of February 2024. These contracts enabled the full-scale operation of the RCGs Secretariat along the year (from March 2023 until Feb 2024). The pan regional ISSG National Correspondents continued the discussions on how to implement the long-term stabilization of the services and tools and ran several consultations with the MS. A main conclusion from these is the fact that the administrative procedures at national level are a continuous and complex constraint for some of the MS. Further discussions on the long-term funding scenarios are planned during 2024. The objective is to develop and implement the long-

term funding solution for the RCGs Secretariat and the tools, while simultaneously, a suitable approach needs to be found to keep the services and tools working continuously to guarantee the support to the RCGs and the LM.

(max. 900 words per study)

SECTION 2: BIOLOGICAL DATA

Text Box 2.1: List of required species/stocks

(Region/RFMO/RFO/IO: Please indicate per text box and update the table of contents)

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.1.

Region: All

Deviations from the work plan

List the deviations (if any) in the achieved data collection (lengths only) compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.1.

In the NAFO area no fish were measured because Estonian vessels did not carry out commercial fishing.

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

There is no solution to avoid deviation in NAFO area. Census approach is used and all the vessels fishing in area have scientific observer onboard collecting samples, but it is not possible to sample commercial catch if commercial fishing is not taking place. Work plans prediction has been based on the reference period fishing effort, but if vessels are not going out then living up to the prediction is not feasible.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.2: Planning of sampling for biological variables

(Region/RFMO/RFO/IO: Please indicate per text box and update the table of contents)

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(a) of the EU MAP Delegated Decision annex. This text box applies to the annual report and complements Table 2.2.

Region: Baltic Sea

Deviations from the work plan

List the deviations (if any) in the achieved collection of biological data (other than lengths), compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.2.

There was one case of under sampling of sex and maturity data because of the declined pikeperch stock compared with the reference period and due to the limitation of fish handling. Some oversampling occurred because of the changed fishing pattern, but according to sampling manual the whole fishing period had to be covered.

Oversampling of sea trout and maturity of salmon was related with higher catches among recruited self sampling fishermen and additionally collected data. Planned salmon maturity number was smaller, because maturity can't be always determined due to the limitations to fish handling.

Actions to avoid deviations.

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

Inventories of collected data are introduced bimonthly to be able to avoid some of these deviations. Inventories will produce an early effect in case there is the possibility to change the sampling intensity.

End user: NAFO

Deviations from the work plan

List the deviations (if any) in the achieved collection of biological data (other than lengths), compared to what was planned.

The general reasons for deviations from the work plan in terms of planned vs. achieved data collection should be summarised in this section, while detailed comments on deviations on particular species/stocks should be included in the 'AR comments' column in Table 2.2.

No samples were taken because Estonian vessels did not carry out commercial fishing in the NAFO area in 2023.

Actions to avoid deviations.

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

There is no solution to avoid deviation in NAFO area. Census approach is used and all the vessels fishing in area have scientific observer onboard taking samples, but it is not possible to sample commercial catch if commercial fishing is not taking place. Work plans

prediction has been based on the reference period fishing effort, but if vessels are not going out then living up to the prediction is not feasible.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.3: Diadromous species data collection in freshwater

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.1(b) and point 2.3 of the EU MAP Delegated Decision annex. Use this text box to give an overview of the methodology used to collect data from freshwater and inland commercial and recreational fisheries for salmon, sea trout and eel. Also include overview of data to be collected from research surveys on salmon, sea trout and eel in freshwater, and on eel in any relevant habitat including coastal waters.

Method selected for collecting data.

Sampling scheme identifier: parrelectrofishing, smolttrapping, adultcounting

Salmon and sea trout

There is no commercial fishery for salmon and sea trout in fresh waters.

The principal way of monitoring salmonid populations in rivers is electrofishing.

Permanent monitoring sites, located in important parr rearing areas, are fished annually or biennially. The sites are fished twice to calculate fishing efficiency and parr densities are presented as individuals per 100 m². The results of this method are comparable to all neighbouring countries. Caught fish are measured and released after analyses.

Atlantic salmon and sea trout smolt abundance estimate in river Pirita is done by capture-mark-recapture method. Smolts are caught by trap-net at the river mouth throughout the migration season. Captured smolts are tagged by VIE (visible implant elastomer) and released 2 km upstream from the trap. Some of the tagged smolts will be recaptured during their descent towards the sea. This enables to estimate the overall smolt run size. Caught fish are measured and released after analyses.

Ascending Atlantic salmon and sea trout spawners are counted in river Pirita throughout the migration season. A fish fence (type: resistance board weir) covering the entire width of the river guides fish through an opening that has a fish counter. Vaki Riverwatcher with a camera tunnel is used.

Recreational fishermen need a special licence to fish salmon and sea trout in rivers and are obliged to report the catches (census survey) to the Ministry of the Environment (entered to TEHA database). Obligation is to send in the report within 5 days after the licence is expired and it is impossible to get new fishing licence if reports of expired licences are not delivered. That way we get the total catch from the database TEHA (Science and Recreational Fisheries database) and we also see non-response rate.

Sampling scheme identifier: freshwater_eelfykes_scientific

Eel

Commercial fishery for eel exists in five lakes and is based on stocked eels. The commercial landing of eel in coastal waters of the Baltic Sea is very limited, average

landings in the reference years (2018-2020) was only 1 tonne.

List of trusted fishermen are used for collecting information from freshwater commercial catches. Total length, weight and silvering stage are recorded for the individuals sampled from commercial catches.

Fishery independent study for eel at coastal areas is done during Coastal fish survey. The fyke nets used are 55 cm high with a semi-circular opening and a leader or wing that is 5 m long. They are made of 17 mm mesh in the arm and 10 mm mesh in the crib of yarn quality no. 210/12 in twisted nylon. Minimum of 80 fyke net days are fished in each of the 6 study areas along the Baltic Sea Estonian coast in July and August. Caught eels are measured for length and weight and otoliths for age reading are collected.

Fishery independent survey in inland waters are carried out in Lake Võrtsjärv. Fyke nets (mouth opening <3m, mesh size >38mm in the cod end) are used. Survey lasts usually from May until October, depending on water temperature. Fyke nets are controlled in every 3 days. Length (TL=mm), weight (g), age (from otolith), silvering stage (length of the pectoral fins and eye diameter) and infestation with parasites are recorded.

Annual data of eel restocking is collected with average weight (g) and total number of restocked individuals recorded.

Recreational fishermen need a special licence to fish eel with long line or harpoon gun in lakes where eels are stocked. With licence comes obligation to report the catches (census survey) to the Ministry of the Environment (entered to TEHA database). Obligation is to send in the report within 5 days after the licence is expired and it is impossible to get new fishing licence if reports of expired licences are not delivered. That way we get the total catch from the database TEHA (Science and Recreational Fisheries database) and we also see non-response rate. Catches made by recreational anglers and rod fishermen are based on regular recreational fishery phone survey questionnaire. Phone survey is conducted in every second year. For recreational anglers and rod fishermen, the catch volume and the confidence limits are calculated and documented from phone survey results.

Phone survey report about catches in 2022 is available at: <https://www.agri.ee/ministeerium-uudised-kontakt/uuringud>

“[Eesti harrastuskalapüügi kvantitatiivuurint 2022. a kohta \(2023\)](#)”

(max 250 words per species and area)

Sampling scheme identifier: parrelectrofishing, smolttrapping, adultcounting

Were the planned numbers achieved? Yes/ No

Yes

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

Sampling scheme identifier: freshwater_eelfykes_scientific

Were the planned numbers achieved? Yes/ No

No.

If the answer is No, explain why not, and what measures were taken to avoid non-conformity.

Fishery dependent eel sampling in Lake Võrtsjärv did not achieve sufficient amount of silver eels due to a very low number of silver eels in the commercial catch. Additional effort was put into collecting samples from fisherman, but with no remarkable benefit to raise the number of silver eels.

(max 250 words per species and area)

Text Box 2.4: Recreational Fisheries

General comment: This text box fulfils Article 5(2)(a), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 2.2 of the EU MAP Delegated Decision annex. Use this text box to give an overview of the methodology used to collect data on marine and freshwater recreational catches.

Region: Baltic Sea

Recreational fishery in Estonia consists of 3 major parts:

- 1) Recreational fishermen using commercial gear (gill nets, trap nets, longlines) have an obligation to report the catches to the Ministry of the Environment.
- 2) Recreational fishermen need a licence to fish salmon and sea trout in rivers and are obliged to report the catches to the Ministry of the Environment. (Covered in text box 2.3)
- 3) Anglers and rod fishermen fishing in sea are not covered with compulsory catch reports. Their catches are estimated during phone survey every second year.

The main target species in the recreational fishery covered with compulsory catch reports are perch, pike-perch, flounder, whitefish, sea trout and salmon (among species listed in the Table 2.1).

Catch report system

Recreational fishermen that use commercial gear or are fishing on “salmonid”-rivers (part 1 and part 2) are obliged to report the catches (census survey) to the Ministry of the Environment (entered to TEHA database). Obligation is to send in the report within 5 days after the licence is expired and it is impossible to get new fishing licence if reports of expired licences are not delivered. That way we get the total catch from the database TEHA (Science and Recreational Fisheries database) and we also see non-response rate.

Phone survey

Catches made by recreational anglers and rod fishermen fishing in the sea (part 3) are based on regular recreational fishery phone survey questionnaire. Phone survey is conducted in every second year

Phone survey report about catches in 2022 is available at:

<https://www.agri.ee/ministeerium-uudised-kontakt/uuringud>

“[Eesti harrastuskalapüügi kvantitatiivuuring 2022. a kohta \(2023\)](#)”

For recreational anglers and rod fishermen fishing at sea, the catch volume and the confidence limits are calculated and documented from phone survey results.

(max 900 words per region)

Region: Baltic Sea

Deviations from the work plan

List the deviations (if any) in the achieved data collection, compared to what was planned in the work plan and explain the reasons for the deviations.

No deviations from the work plan.

Phone survey took place at the beginning of 2023 with an aim to collect catch information about 2022. Everything went according to the plan.

Catch reporting when using fishing card is compulsory and we have official data there.

Table 2.4 of this AR includes additional rows on the availability of recreational catch report data for cod, salmon and trout, which were erroneously omitted from the work plan.

Action to avoid deviations

Describe the actions that will be considered/have been taken to avoid the deviations in the future and when these actions are expected to produce results.

(max 900 words per region)

Text Box 2.5: Sampling plan description for biological data
(Region/RFMO/RFO/IO: Please indicate per text box and update the table of contents)

General Comment: This text box fulfils Article 5(2)(a) and (b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter 2 point 2.1(a) of the EU MAP Delegated Decision annex. This text box complements Table 2.5.

This text box is complementary to information on the sampling schemes provided in the quality document (Annex 1.1). It serves to highlight additional information on sampling schemes and sampling frames that the Member State considers useful to understand the sampling design planned for the region and the implementation year(s).

Region: Baltic Sea

Deviations from the work plan

List deviations (if any) in the achieved data collection compared to what was planned in the work plan and explain the reasons for the deviations.

There were less PSUs sampled in small boat coastal fishery because of the declined stocks. Therefore, fishing effort was concentrated to less months and in some months with small fishing effort sampling was not feasible.

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

To achieve planned PSU numbers sampled in SB coastal fishery inventories of collected data were introduced bimonthly to be able to prevent deviations. Inventories will produce an early effect in case there is a possibility to change the sampling intensity.

End user: NAFO

Deviations from the work plan

List deviations (if any) in the achieved data collection compared to what was planned in the work plan and explain the reasons for the deviations.

No samples were taken because Estonian vessels did not carry out commercial fishing in the NAFO area in 2023.

Actions to avoid deviations

Describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

There is no solution to avoid deviation in NAFO area. Census approach is used and all the vessels fishing in area have scientific observer onboard, but it is not possible to sample commercial catch if commercial fishing is not taking place. Work plans prediction has been based on the reference period fishing effort, but if vessels are not going out then living up to the prediction is not feasible.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 2.6: Research surveys at sea

(Research survey: Please indicate per text box and update the table of contents)

General Comment: This Text box fulfils Article 5 (1)(b), Article 6 (3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapters I and II of the EU MAP Implementing Decision. It is intended to specify which research surveys at sea set out in Table 2 of the EU MAP Implementing Decision will be carried out. Member States shall specify whether the research survey is included in Table 2 of the EU-MAP Implementing Decision or whether it is an additional survey.

Region: Baltic Sea

Baltic International Trawl Survey (BITS Q4)

1. Objectives of the survey

Survey is included in Table 2 of the EU-MAP Implementing Decision. Target species are demersal fish species, mainly Baltic cod and flatfish species (flounder, plaice, dab, brill and turbot). The main aim is to determine the year-class strength of the target species. Target data are abundances, weight and length distributions of all fishes and length-weight-age- sex-maturity data of commercially important species as well as hydrographic data (temperature, salinity and oxygen). In addition, cod parasite infection load in livers and marine litter are sampled and marine mammal observations are recorded.

In Estonian survey area species as BLL, DAB and PLE are extremely rare.

The results are used by the ICES WGBFAS for assessment of the Baltic cod and flounder stocks on routinely basis.

Survey takes place in October/November.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in

Table 2.6 for this specific survey.

Estonia follows the methods agreed for the BITS by the ICES Baltic International Fish Survey Working Group (WGBIFS) described in the Manual for the Baltic International Trawl Surveys (BITS).

<http://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20%28SISP%29/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20%28BITS%29.pdf>

3. For internationally coordinated surveys, describe the participating Member States/vessels.

The participating countries use their research vessel or chartered fishing vessel and the standard gear. Estonia is participating in the 4th quarter (autumn) survey using the chartered Estonian fishing vessel and TV2-520 trawl.

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

The ICES survey planning group (WGBIFS) assigns the tasks to the survey participants (e.g. coverage of certain areas in a certain time frame). Each participating country is responsible for the activities conducted on its national part of the international survey. 10 fish hauls is maximum expected number for Estonia, however the actual number may differ by year.

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

ICES Scientific Reports. 5:70. 90 pp. ICES Scientific Reports. 5:70. 90 pp.

<https://doi.org/10.17895/ices.pub.23675049>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

The Baltic International Survey (BITS) survey results are used to estimate stock indices of the commercially exploited targeted fish species in the Baltic mainly Baltic cod and the flatfish species flounder, plaice, dab, brill and turbot. This long time series of fisheries-independent survey provides the data for fisheries assessments and the sustainable

management of Baltic fish stocks in the frame of ICES WG's, mainly WGBFAS.

7. Extended comments

Extended AR comments can be placed under this section.

5 trawl hauls allocated for Estonia were performed in the ICES Sub-division 28.2. The number and location of hauls to be executed by the MS has been defined by the ICES WGBIFS.

(max. 450 words per survey)

Baltic International Acoustic Survey (BIAS)

1. Objectives of the survey

Survey is included in Table 2 of the EU-MAP Implementing Decision. Target species of the Baltic International Acoustic Survey (BIAS) are small pelagic fish species, mainly Baltic herring and sprat. The main aim is to provide information on stock parameters of small pelagics in the Baltic Sea. Target data are biomass, weight and length distributions and length-weight-age-sex-maturity of small pelagics, as well as hydrographic data (temperature, salinity and oxygen).

Survey takes place in September/October.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Baltic International Acoustic Survey (BIAS) is carried out in September/October. Surveys is carried out according to the agreed Manual of International Baltic Acoustic Surveys (IBAS):


[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Each MS performs the survey in its EEZ on its own or shared research vessel. Estonia is chartering the Polish Research vessel BALTICA. The overall coordination of the surveys is done by the WGBIFS in order to secure the full coverage of the Baltic Sea.

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

The ICES survey planning group (WGBIFS) assigns the tasks to the survey participants (e.g. coverage of certain areas in a certain time frame). Each participating country is responsible for the activities conducted on its national part of the international survey.



Map: Baltic International Acoustic Survey (BIAS), Sep/Oct 2018: Survey areas by country.

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

ICES Scientific Reports. 5:70. 90 pp. <https://doi.org/10.17895/ices.pub.23675049>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context.

If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.

BIAS survey provide important fishery-independent estimates for Baltic herring and sprat stocks and the results are used by the ICES WGBFAS for assessment of the Baltic herring and sprat stocks on routinely basis.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Gulf of Riga Acoustic Herring Survey GRAHS

1. Objectives of the survey

Survey is included in Table 2 of the EU-MAP Implementing Decision. The aim of the survey is to obtain the fisheries-independent information for tuning analytical stock assessment models for Baltic herring in the Gulf of Riga (Gulf of Riga herring). Target data are biomass, weight and length distributions and length-weight-age-sex-maturity of Baltic herring, as well as hydrographic data (temperature, salinity and oxygen). The information obtained during the survey is used by the Baltic Fisheries Assessment Working Group of the ICES (WGBFAS).

Survey takes place in July/August.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Survey will be carried out following the agreed Manual of International Baltic Acoustic Surveys (IBAS)

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

The surveys are coordinated and the results are discussed by the ICES WGBIFS annually. The survey is carried out in July-August annually in order to cover the period after main spawning season when most of the stock has left the near-coast spawning grounds.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

The survey is carried out jointly by the Latvian and Estonian scientists on the chartered Latvian fishing vessel.

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

Estonia and Latvia share the tasks of work and also the survey costs on this joint survey.

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

ICES Scientific Reports. 5:70. 90 pp. <https://doi.org/10.17895/ices.pub.23675049>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

GRAHS survey provide important fishery-independent estimates for Gulf of Riga herring stock and the results are used by the ICES WGBFAS for assessment of the Baltic herring stock on routinely basis.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Sprat Acoustic Survey (SPRAS)

1. Objectives of the survey

Survey is included in Table 2 of the EU-MAP Implementing Decision. Target species is sprat. The main aim is to provide information on stock parameters of sprat in the Baltic Sea. Target data are biomass, weight and length distributions and length-weight-age-sex- maturity of sprat as well as hydrographic data (temperature, salinity and oxygen).

Survey takes place in May.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

The Baltic Acoustic Spring Survey (SPRAS) is carried out annually in May. Survey is carried out by the agreed Manual of International Baltic Acoustic Surveys (IBAS) ([https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)).

The survey is annually coordinated and the results discussed by the ICES WGBIFS.

The acoustic survey covers the main area of sprat distribution in the Baltic. Each statistical rectangle of the area under investigation is allocated to one particular country by the Baltic International Fish Survey Working Group (WGBIFS), thus each country has a mandatory responsible area.

3. For internationally coordinated surveys, describe the participating Member States/vessels.

Each MS performs the survey in its EEZ on its own or shared research vessel. Estonia is chartering the Polish Research vessel BALTICA for SPRAS surveys. The overall

coordination of the coming surveys is done by the WGBIFS in order to secure the full coverage of the Baltic Sea.

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

The ICES survey planning group (WGBIFS) assigns the tasks to the survey participants (e.g. coverage of certain areas in a certain time frame). Each participating country is responsible for the activities conducted on its national part of the international survey.

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

ICES Scientific Reports. 5:70. 90 pp. <https://doi.org/10.17895/ices.pub.23675049>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

SPARS survey provide important fishery-independent estimates for the Baltic Sea sprat stock and the results are used by the ICES WGBFAS for assessment of the sprat stock on routinely basis.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Estonian Fish Larvae Survey

1. Objectives of the survey

This national survey has been conducted annually since 1947 to study the distribution and abundance of fish larvae and juveniles with the aim to provide primary information on herring yearclass abundance and to understand the affecting environmental background. Results of the survey is used for early estimation of the spawning success and yearclass potential of herring, but also commercially important percids (perch and pikeperch) and smelt. Data on abundance and distribution of the spring spawning herring larvae in the Gulf of Riga is used to calculate larval herring index N2. The rationale for this approach is that (i) the eggs and larvae of a species should be in direct proportion to the biomass of the adults, (ii) estimating the abundance of eggs or larvae is either more accurate or less costly than estimating the abundance of adults, and (iii) ichthyoplankton surveys provide data on a

different group of species than trawl surveys for adults. Data of the survey is used nationally for Baltic herring and smelt stock assessment and for assessment of the state of important spawning grounds.

Survey takes place in May till July.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Nine fixed stations are visited weekly (May-July). Hensen larval fish net is used for 10 min. hauls in NE of the Gulf of Riga in commercially important fish spawning and nursery grounds.

Larvae and juveniles are collected using research vessels of the Estonian Marine Institute. Larvae are identified, measured and counted. Plankton samples and environmental data are collected and analysed.

Details of the survey design is described in following paper:

<https://www.sciencedirect.com/science/article/abs/pii/S016578361000281X>

Mesozooplankton samples are collected according to HELCOM manual:

<https://helcom.fi/media/publications/Guidelines-for-monitoring-of-mesozooplankton.pdf>

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

<https://www.agri.ee/media/11646/download>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

Data of the survey is used nationally for Baltic herring and smelt stock assessment and for assessment of the state of the spawning grounds. Larval herring index N2 is calculated.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Gulf of Riga Fish survey

1. Objectives of the survey

Objective of this national survey is to collect fisheries independent data for tuning the pikeperch and perch VPA assessment and get abundance index and biological data about other commercially important species (smelt, cyprinids).

Trawl survey in the Pärnu Bay was conducted already in earlier decades (since 1950s) and resumed 2009. Data of the survey is used nationally for stock assessment of many coastal fish species including perch and pikeperch.

Survey takes place in May and September till December.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Research vessel of the Estonian Marine Institute AURELIE is used for trawling. The trawl (working depth 0.3 m from the bottom) is pulled with the speed of 3 knots for 30 minutes. The trawl mouth is 2 m high and 6 m wide, distance between doors is 20 m and maximum distance between the 8.2 m long trawl wings is 12 m. Mesh size is 60 mm (knot to knot), at the tip of the trawl wings, 45 mm at the trawl mouth and decreases gradually to 10 mm at the codend. Six fixed trawl transects are situated three to eight km from shore (water depth five to nine m) to cover the entire length of the Pärnu Bay. All fixed stations are fished in May, September, October, November and December. The survey is coordinated and the results are discussed by the HELCOM FISH-PRO annually. Survey is carried out following the manual of HELCOM: (<https://portal.helcom.fi/meetings/FISH-PRO%20III%201-2019-592/MeetingDocuments/4-1%20Updated%20guidelines%20for%20coastal%20fish%20monitoring.pdf>)

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any

status report (e.g. Cruise report).

<https://www.agri.ee/media/11635/download>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

Data of the survey is used nationally for stock assessment of many coastal fish species including for calculating fishery-independent estimates for the analytical assessment of the perch and pikeperch stocks.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

Coastal fish survey

1. Objectives of the survey

National annual survey started in 1992, now in 10 fixed areas. Data on abundance, relative year class strength, size and age distribution, structure of coastal fish assemblages are used in routine basis for giving advice for main commercially important species (also those not namely listed in the National WP). Most of our coastal fish are listed in Commission Delegated decision (EU) 2021/1167 Table 2 as relevant to HELCOM. Additional data on bycatch (incl. birds), seal damage to catch, distribution and abundance of alien species, microplastics in fish stomachs is collected. Data from the survey is used to calculate HELCOM and Marine Strategy Framework Directive fish indicators. Collected information (CPUE, age and length distribution, age-length keys for commercial species etc) form the basis for advice for commercially important stocks (perch, pikeperch, flounder, cyprinids) including fisheries independent data for tuning the flounder, pikeperch and perch analytical assessment.

Survey takes place in May till August, in October and in November.

2. Description of the survey design and methods used in the survey for each type of data collection as listed in Table 2.6 for this specific survey.

Annual gill-net survey in defined areas along the Estonian coast, in fixed (Pärnu, Hiiumaa, Kõiguste and Küdema) or random stations (Narva-Jõesuu, Vilsandi, Kihnu, Matsalu, Käsmu, Vaindlo). Each station consists of a series of gill nets of fixed mesh sizes and construction. 6-72 stations are fished in each area. Additionally in six areas (Vilsandi, Kihnu, Kõiguste, Matsalu, Saarnaki and Käsmu) minimum 80 standard fyke net/days are fished for eel CPUE and length distribution.

Methods: Thoresson, 1995, HELCOM 2015

(<http://www.helcom.fi/Documents/Action%20areas/Monitoring%20and%20assessment/Ma>

nuals% 20and% 20Guidelines/Guidelines% 20for% 20Coastal% 20fish% 20Monitoring% 20of
% 20HELCOM.pdf)

3. For internationally coordinated surveys, describe the participating Member States/vessels.

N/A

4. Where applicable, provide more details on the type of participation and/or threshold agreement applied.

N/A

(max 450 words per survey)

5. For internationally coordinated surveys, provide a link to the latest meeting report of the coordination group.

Provide a link to the meeting report from the body coordinating the survey (ICES, MEDITS coordination group, MEDIAS coordination group, etc.). For surveys that are not internationally coordinated, refer to any status report (e.g. Cruise report).

<https://www.agri.ee/media/11635/download>

6. List the main use of the results of the survey (e.g. indices, abundance estimates, environmental indicators). Specify in which context the results are used (on a routine basis), both in international and national context. *If presenting maps of the achieved research survey stations is necessary, provide them as an annex. Refer clearly to the annex and map numbers.*

Data from the survey is used to calculate HELCOM and Marine Strategy Framework Directive fish indicators. Collected information (CPUE, age and length distribution, age-length keys for commercial species etc) form the basis for advice for commercially important stocks (perch, pikeperch, flounder, cyprinids) including fisheries independent data for tuning the flounder and perch analytical assessment.

7. Extended comments

Extended AR comments can be placed under this section.

(max. 450 words per survey)

SECTION 3: FISHING ACTIVITY DATA

Text Box 3.2: Fishing activity variables data collection strategy (for inland eel commercial fisheries)

General comment: This text box fulfils Article 5(2)(c), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter II point 3.2 of the EU MAP Delegated Decision annex. It is intended to describe the methods and data sources used to estimate fishing capacity, effort and landings data.

Freshwater eel fishery (10-35 t/year, 2006-2020) – occurs in Narva RBD. All of the eel caught is of restocked background. The gear used consists mostly of fyke nets, very seldom long lines are used. The total capacity of the freshwater fishery in 2020 was 359 commercial fishermen/companies. 85 commercial fishermen/companies of the freshwater fishery reported eel in their catch. In the freshwater fishery 96% (35.8 t) of the eel was caught from Lake Võrtsjärv by 48 fisher-men/companies (averaging 745.1 kg per company/fisherman, CPUE per gear/night = 0.8 kg). This information is collected by the Estonian Ministry of Rural Affairs. Register is updated every year and is available online. Records are kept over the number and type of gears used. Data from fishermen logbooks are collected once a month and uploaded twice a year.

(max. 900 words)

Deviations from the work plan

List the changes from the work plan (if any) and explain the reasons.

In 2021, 2022 and 2023 the eel catches from Narva RBD have significantly increased to 46, 50,8 and 57,1 t/year, respectively. As Commercial fishing information is collected by the Republic of Estonia Agriculture and Food Board.

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

(max. 900 words)

SECTION 4: IMPACT OF FISHERIES ON MARINE BIOLOGICAL RESOURCES

Text Box 4.2: Incidental catches of sensitive species

(Region/RFMO/RFO/IO: Please indicate per text box and update the table of contents)

General Comment: This text box fulfils Article 5(2)(a) and (b), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004 and Chapter 2 point 4.1 of the EU-MAP Delegated Decision annex. This text box complements Table 2.5.

Region: Baltic Sea

Additional information on planning the observation of incidental catches of sensitive species (if already filled in in Annex 1.1, please indicate where it can be found):

- Has an assessment of the relative risk of bycatch for the different gear types/metiers taken place and been taken into account for the sampling design?

Yes.

- What are the gear types/metiers that present the highest risk of bycatch per species/taxa of PETS in a given region?

Gill nets present the highest risk of bycatch to diving waterbirds and fyke nets are the risk to the seals in the Baltic Sea.

Baltic pelagic trawlers, targeting sprat and herring, are considered as low risk for PETS bycatch.

- What are the methods to calculate the observation effort?

There is census approach: registering bycatch is mandatory by the law for all fishermen and data are included in the Estonian Fisheries Information System. Addition to that we carry out direct questioning of the crew to collect information about sighting/catching whales (*Phocoena phocoena*) in our Baltic trawl fisheries (exclusively pelagic trawls for sprat and herring). This coverage is the same as for sampling schemes identified as OnShoreCommercialPelagic and OnShoreCommercialPelagicGOR. For our long distance fleet we use on-board observers to collect data on any bycatch. In NAFO area on-board observer coverage is 100% and in NEAFC the coverage has been between 10 to 20%.

Bycatch information is recorded during all surveys, including our rather large scale gill net survey (Coastal Fish Survey).

Recreational fishermen fishing with commercial gear are also obliged to report any occurred bycatch.

- Does the sampling design and protocol follow the recommendations from relevant expert groups If there are no relevant expert groups, the design and protocol have to be explained in the text?

Sampling design and protocol does not follow the recommendations from relevant expert groups, although ICES WGBYC reports are monitored. Planned number of PSUs for trawls (OSF PEL, GOR PEL) in Table 2.5 cover the direct questioning of the crew about seeing or catching the whales (*Phocoena phocoena*). Trips are checked and the crew is questioned in ports by the staff of the EMI-UT during taking the herring and sprat samples. No mammal nor bird bycatch have been registered since the beginning of the commitment.

Observers covering our long distance fleet have an obligation to monitor hauling on deck to notice and register any bycatch event. It is also noted down when observer was not present during the hauling.

Additional information on observer protocols (if already filled in in Annex 1.1, indicate where it can be found):

- Does the on-board observer protocol contain a check for rare specimens in the catch at opening of the cod-end? If YES is the observer instructed to indicate if the cod-end was NOT checked in a haul?

Yes and if the codend was not checked in a haul, observers should indicate it in the report. Observer is instructed to report all bycatch in haul, not only mammals or birds or sensitive species.

- In gill nets - and hook-and-line fisheries: does the on-board observer protocol instruct the observer to indicate how much of the hauling process has been observed for (large) incidental bycatches that slip out of the net?

There is no dedicated observer programme for gill net nor hook-and-line fishery in Estonia. Data are collected through log books. In gill net fishery only small boats (below 12m) are used and it is seldom possible to put observers on board for safety reasons. Using CCTV or any other similar approach is not yet possible due to lack of national legislation that would allow it. Work on changing the legislation is ongoing, but progress has been rather slow due

to strong resistance from different groups. Hook-and-line is seldom used nowadays as main species fished by this technique was eel.

All fishermen are obliged to report all bycatches and all alive specimens of incidental catch should be released.

- In large catches: does the protocol instruct to check for rare specimens during sorting of the catch (i.e. at conveyor belt)? Is the observer instructed to indicate what percentage of the sorting or hauling process has been checked at "haul level"?

Y, Y.

Additional information on sampling schemes

Pilot study about bird and seal bycatch in our coastal fishery is ongoing (conducted during the period 2020-2021). Results of the pilot study will be compared to a similar study conducted in 2007-2009 (part of a LIFE project). Based on this study results and analyses a decision will be made on how often this kind of bycatch study should be conducted in the future.

Additional description on sampling frames

(One text box (max. 1 000 words) per region/RFMO/RFO/IO)

Region: Baltic Sea

Results

Provide additional information, if available, in this text box. For example, summary information on the number of individuals recorded as bycaught per species, gear group and monitoring method with information about the state of the animals (i.e. were they released alive, dead, or collected for sampling).

In 2023 by-catch incidents happened in coastal fishery, but not in trawl fishery. In the coastal fishery (total of 30 509 fishing trips) by-catch was registered in 71 cases: mammals in 31 cases (altogether 45 grey seals (*Halichoerus grypus*) and 4 Eurasian otters (*Lutra lutra*) and birds in 40 cases (in total 108 specimens: unidentified ducks 24, 68 great cormorants (*Phalacrocorax carbo*), 4 common mergansers (*Mergus merganser*), 1 tufted duck (*Aythya fuligula*), 1 great crested grebe (*Podiceps cristatus*), 1 wadder *Tringa sp.*, 8 long-tailed ducks (*Clangula hyemalis*) and 1 unidentified bird. Seal damage to fishing gear or catch was registered in 399 cases and Eurasian otter damage in 3 cases. Cormorant damage to catch or gear was reported in 26 cases. During recreational fishery with gillnets 32 bird bycatch incidents were recorded.

Test study: Accidental bycatch of birds and seals in passive commercial fishing gear of Estonian coastal fishery 2020-2021

Data on bycatch of seals and birds from passive commercial fishing gear (fykes and gillnets) was collected by contracting commercial coastal fishers to register and report data about all the bycatch occasions during 2020 and 2021. At the same time also data on fishing effort was registered so catch per unit effort of bycatch and estimate of bycaught individuals over all coastal fishery were calculated for 2021. Data were received from 39 commercial fishers from all Estonian coastal areas (Gulf of Finland, Väinameri, Gulf of Riga and Baltic Proper). 248 bycatch cases were registered. Bycatch of four mammal (grey seal, ringed seal, Eurasian beaver and Eurasian otter) and 15 bird species were registered. Grey seal was the most abundant mammal species registered as accidental bycatch in

fishing gear was (52 registered in study, estimate of 110±22 individuals caught in 2021). Other mammals were registered as single individuals (one beaver and one otter) and two ringed seals caught allowed for estimated bycatch of this species to 20 (±10) individuals in Estonian coastal areas in 2021. From 59 bycatch cases of mammals 52 individuals were caught with fykes and 7 with gillnets. In case of birds, the most abundant species caught in passive fishing gear was great cormorant (65 registered in study, estimate of 286±48 individuals caught in 2021). From 139 bycatch cases of birds 68 were caught in fykes and 71 in gillnets. In comparison to previous analogous studies from Estonian coastal fishery, this study demonstrated considerably lower bycatch estimates of seals (estimated annual bycatch of 260-370 individuals in Vetemaa 2009 and 780-930 individuals in Vanhatalo et al. 2014). Also, Vetemaa (2009) estimated 2154 birds to be caught in passive fishing gear in Estonian coastal sea annually – a considerably higher estimate than registered in this study. These differences may occur due to methodological differences in estimate calculations but may also reflect changes in fishing practises and gear used (e.g. modification of fykes to avoid bycatch of seals). The results of current study also indicate that species abundance in the area is positively associated with species abundance in bycatch. This is illustrated by replacement of long tailed duck as the most numerous bird caught in fishing gear (Vetemaa 2009) with great cormorant in current study. Current study also revealed that 16,6% of birds that were caught in passive gear were released unharmed. Therefore, this study indicates that in the case of most species that were registered as bycatch in commercial fishing gear in current study, the mortality due to bycatch in fishing gear has probably very small effect on population level.

The study is available at: <https://www.agri.ee/media/9888/download>

To estimate the unreported bycatch, it has been decided to proceed with data collection by selected contracted commercial coastal fishermen to register and report data about accidental bycatch of birds and seals in passive commercial fishing gear in coastal fishery.

Deviations from the work plan

The Member State shall list the deviations (if any) in the achieved data collection compared to what was planned in the work plan and explain the reasons for the deviations.

No deviations from the work plan.

Actions to avoid deviations

The Member State shall describe the actions that will be considered/have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

(One text box of max. 1 000 words per region/RFMO/RFO/IO)

Text Box 4.3: Fisheries impact on marine habitats

General comment: This text box fulfils Article 5 paragraph 2(a) and 2(b), Article 6 paragraph 3(a), 3(b) and 3(c) of Regulation (EU) 2017/1004 and Chapter 2, section 4.2 of the EU MAP Delegated Decision annex. It contains information on additional studies on the fisheries impact on marine habitats.

At the moment no additional studies are planned to address the fisheries impact on marine habitats. As there are no active bottom trawlers in our Baltic Sea fleet, so we do not study fishery's impact to the sea floor.

(max 900 words per study)

Brief description of the results (including deviations from the plan and justifications as to why if this was the case).

Achievement of the original expected outcomes and justification if this was not the case.

NA

Follow-up to the activities (what are the next steps, how the results will be used).

NA

(max. 900 words per study)

SECTION 5: ECONOMIC AND SOCIAL DATA IN FISHERIES

Text Box 5.2: Economic and social variables for fisheries data collection

General comment: This Text box fulfils Article 5(2)(d), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004, and Chapter II point 5 of the EU MAP Delegated Decision annex. It is intended to specify data to be collected under Tables 7, 8 and 9 of the EU MAP Delegated Decision annex.

1. Description of clustering

Estonia has two clustered fleet segments. In both cases the segments added were with same fishing techniques and were merged for confidentiality reasons. Clusters were named after the biggest segment in terms of number of vessels.

Clustered fleet segments:

- 1) Pelagic trawlers 24-<40 m* – this segment catches sprat and herring in the Baltic Sea. Vessels of three length classes are merged. The number of vessels in each length class in 2020 was as follows:
 - a. 12-<18 m one vessel,
 - b. 18-<24 m six vessels and
 - c. 24-<40 m 20 vessels.
- 2) Demersal trawlers and/or demersal seiners 40 m or larger* – this segment operates in the NAFO and NEAFC areas. Vessels of two length classes are merged. The number of vessels in both length classes in 2020 was as follows:
 - a. 24-<40 m one vessel and
 - b. 40 m or larger five vessels.

2. Description of activity indicator

Estonia does not use the activity indicator

3. Deviation from the RCG ECON (ex. PGECON) definitions

No deviations are planned.

(max. 900 words)

Deviations from the work plan

Estonia does not use the PIM. The data for capital value and capital costs is obtained from financial accounts of fishing companies. The EU MAP Guidance Document for the Fishing Fleet states that MS may use ‘Alternative methods based on company surveys. These alternative methods may be used if the derived estimates reflect the actual definition of net capital stock (depreciated replacement value of the vessel including on-board equipment with a useful lifetime of more than one year)’.

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

(max. 900 words)

SECTION 6: ECONOMIC AND SOCIAL DATA IN AQUACULTURE

Text Box 6.1: Economic and social variables for aquaculture data collection

General comment: This text box fulfils Article 5(2)(e), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004, and Chapter II point 6 of the EU MAP Delegated Decision annex. It is intended to specify data to be collected under Tables 10 and 11 of the EU MAP Delegated Decision annex.

Estonia will not collect data on the aquaculture. As established by Commission Implementing Decision (EU) 2021/1168 it is not mandatory for Member States to collect social, economic and environmental data on aquaculture if their total aquaculture production is less than 1 % of total Union aquaculture production. According to the latest EUROSTAT data from 2019 the Estonian aquaculture production (1062 t) was below the threshold, accounting for 0,001% of the total aquaculture production of EU in 2019.

(max. 900 words)

Deviations from the work plan

NA

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

(max. 900 words)

SECTION 7: ECONOMIC AND SOCIAL DATA IN FISH PROCESSING

Text Box 7.1: Economic and social variables for fish processing data collection

General comment: This text box fulfils Article 5(2)(f), Article 6(3)(a), (b) and (c) of Regulation (EU) 2017/1004, and Chapter II point 7 of the EU MAP Delegated Decision annex.

According to the EU MAP the collection of data on the processing industry is optional. Because similar data is already collected by Statistics Estonia and forwarded to EUROSTAT, Estonia will not collect data on the processing industry to avoid the duplication in data collection.

(max. 900 words)

Deviations from the work plan

NA

Actions to avoid deviations

Briefly describe the actions that will be considered / have been taken to avoid deviations in the future and when these actions are expected to produce an effect. If there are no deviations, then this section is not applicable.

(max. 900 words)

ANNEX 1.1 - QUALITY REPORT FOR BIOLOGICAL DATA SAMPLING SCHEME

The quality report fulfils Article 6(3)(d) of Regulation (EU) 2017/1004. This document is intended to specify data to be collected under Chapter II, point 2 of the EU MAP Delegated Decision annex: Biological data on exploited biological resources caught by Union commercial and recreational fisheries. Use this document to state whether documentation in the data collection process (design, sampling implementation, data capture, data storage, sample storage and data processing) exists and identify where this documentation can be found. Names of sampling schemes and strata shall be identical to those in Tables 2.2, 2.3, 2.4, 2.5, 2.6 and 4.1 of the WP/AR. In case of quality information on scientific surveys, use the survey acronym as a sampling scheme identifier. For mandatory surveys, refer to Table 1 of the EU MAP Implementing Decision annex, see also MasterCodeList 'Mandatory survey at sea'.

OnShoreCommercialCoastal

MS: EST
Region: Baltic Sea
Sampling scheme identifier: OnShoreCommercialCoastal
Sampling scheme type: Commercial fishing trip
Observation type: SciObsOnShore
Time period of validity: 2022-2027
Short description (max 100 words): Sampling scheme aiming at collecting biological samples from commercial landings on-shore about selected species in costal fishery using passive gears and small boats (<12m). The scheme covers areas SD28-32.
Description of the population
Population targeted: PSU is month and sampling is species specific and evenly distributed by catch volumes along the coastal area. In total 4 species are sampled in different sub areas: <i>Platichthys flesus</i> (SD28-29, 32), <i>Perca fluviatilis</i> (SD28.1), <i>Sander lucioperca</i> (SD28.1) and <i>Clupea harengus</i> (caught with FPN during spawning season in SD28.1, 29, 32).
Population sampled: Active fishing months when relevant species are fished are covered. Some months with no or very low catches are excluded.
Stratification: <i>Platichthys flesus</i> in SD29 and SD32 and <i>Platichthys flesus</i> in SD28, and <i>Clupea harengus</i> in SD28.1 and <i>Clupea harengus</i> in SD29 and 32 are stratified because belonging to the different management units.
Sampling design and protocols

Sampling design description: Method of PSU selection is NPJS. Selection of month takes place based on the individual expert judgement taking into account the fishing effort and catch volume of selected coastal fish species. Sampling scheme follow the Hierarchy 8 in RDBES: sampling time (month) is the primary sampling unit (PSU) and vessel is the secondary sampling unit (SSU). Sampling scheme applies to all fractions, excl. PETS.

Is the sampling design compliant with the 4S principle?: N

Regional coordination: N

Link to sampling design documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: Y

Monitoring of sampling progress within the sampling year: On weekly basis online catch and landings reports are used to evaluate the fishing activity in different areas and quota exhaustion. This data is compared against the number of samples collected to assure the implementation of sampling program.

Data capture

Means of data capture: Measuring board, scales, tools to collect age reading structures like operculums and otoliths.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: In aggregated form, the data is transmitted to Intercatch and the detailed data to Fishframe or RDB database.

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: Thoresson, G. 1996. Guidelines for coastal monitoring. Kustrapport 1: 1-35. <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx> Estonia uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length scatter plots by visual inspection.

AR comment: Indicate any deviations or developments. Do not change the text already adopted in the work plan.

Less PSUs sampled (83%) because of the declined catches of all relevant coastal species (*Platichthys flesus*, *Clupea harengus*, *Perca fluviatilis* and *Sander lucioperca*).

Sampling design documents in English can be found:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

SelfOnShoreCommercialCoastalSalmon

MS: EST

Region: Baltic Sea

Sampling scheme identifier: SelfOnShoreCommercialCoastalSalmon

Sampling scheme type: Commercial fishing trip

Observation type: SelfOnShore

Time period of validity: 2022-2027

Short description (max 100 words): self-sampling scheme aiming at collecting biological samples from commercial landings on-shore about *Salmo salar* and *Salmo trutta* in costal fishery using passive gears and small boats (<12m). The scheme covers areas SD28-32.

Description of the population

Population targeted: PSU is month and sampling is species specific and covers relevant areas. In total 2 species are sampled in different sub areas: *Salmo salar* (SD22-31 and 32) and *Salmo trutta* (SD22-32).

Population sampled: active fishing months when relevant species are fished are covered. Some months with no or very low catches are excluded.

Stratification: *Salmo salar* in SD22-31 and SD32 is stratified because belonging to the different management units.

Sampling design and protocols

Sampling design description: Method of PSU selection is NPJS. Selection of month takes place based on the individual expert judgement taking into account the fishing effort and catch volume of selected fish species. Sampling scheme follow the Hierarchy 8 in RDBES: sampling time (month) is the primary sampling unit (PSU) and vessel is the secondary sampling unit (SSU). Sampling scheme applies to all fractions, excl. PETS.

Is the sampling design compliant with the 4S principle?: N

Regional coordination: N

Link to sampling design documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Sampling implementation

<p>Recording of refusal rate: Y</p>
<p>Monitoring of sampling progress within the sampling year: Recruited fishermen are interviewed on weekly or monthly bases about their success to collect the samples. If sampling program is hindered by low numbers more fishermen are recruited.</p>
<p>Data capture</p>
<p>Means of data capture: Measuring board, scales, tools to collect age reading structures like operculums and otoliths.</p>
<p>Data capture documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Quality checks documentation: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Data storage</p>
<p>National database: EMI-UT</p>
<p>International database: In aggregated form, the data is transmitted to Intercatch and the detailed data to Fishframe or RDB database.</p>
<p>Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Sample storage</p>
<p>Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.</p>
<p>Sample analysis: Estonia takes part in intercalibration workshops. https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx</p>
<p>Data processing</p>

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length scatter plots by visual inspection.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Sampling design documents in English can be found:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion-methodologi>

OnShoreCommercialPelagic

MS: EST

Region: Baltic Sea

Sampling scheme identifier: OnShoreCommercialPelagic

Sampling scheme type: Commercial fishing trip

Observation type: SciObsOnShore

Time period of validity: 2022-2027

Short description (max 100 words): sampling scheme aiming at collecting biological samples from commercial landings on-shore about *Clupea harengus* and *Sprattus sprattus* in Baltic Sea pelagic trawlers (vessels 12-40m boats). The scheme covers areas SD25-29, 32.

Description of the population

Population targeted: PSU is week. Average number of PSU during the reference years was calculated as the sum of subareas. E.g. the number of weeks in 2020 when fishing was conducted was as follows: SD 28.2 - 34 weeks; SD29 - 37 weeks, SD32 - 44 weeks. The total number of weeks in 2020 when fishing was ongoing is calculated as the sum of weeks in all subareas (34+37+44=115). Same is done for all reference years and then divided by 3 to get the average per reference time period.

Population sampled: On average 2 weeks per month per area are sampled during the active fishing period. No weeks are excluded during the planning.

Stratification: The fishing area consists of multiple subareas (SD29, 28.2, 32) and each of these is sampled separately, as the aim is to get detailed description of the fisheries per subarea, even though all these subareas belong under one stock unit (Central Baltic herring stock and Baltic sprat stock unit)

Sampling design and protocols

Sampling design description: Primary sampling unit (PSU) is time unit of week. Method of PSU selection is NPJS. Sampling scheme for these segments follow the Hierarchy 8 in RDBES: sampling time (weeks) is the primary sampling unit (PSU) and vessel is the secondary sampling units (SSU). For the pelagic fisheries (OSF PEL, GOR PEL) our sampling strategy is that in every month when active fishing is ongoing (this depends on the year and subarea) subset of week(s) are sampled (PSU=week). Sampling scheme applies to all fractions (crews are interviewed about the presence of whales, while collecting the fish sample).

Is the sampling design compliant with the 4S principle?: N

Regional coordination: Y, Multi-lateral agreement with Finland, see table 1.3.

Link to sampling design documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: Y

Monitoring of sampling progress within the sampling year: On weekly basis online catch and landings reports are used to evaluate the fishing activity in different areas and quota exhaustion. This data is compared against the number of samples collected to assure the implementation of the sampling program.

Data capture

Means of data capture: Measuring board, scales, tools to collect otoliths.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: In aggregated form, the data is transmitted to Intercatch and the detailed data to Fishframe or RDB database.

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx>
EMI-UT uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length scatter plots by visual inspection.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. Sampling design documents in English can be found:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

OnShoreCommercialPelagicGOR

MS: EST

Region: Baltic Sea

Sampling scheme identifier: OnShoreCommercialPelagicGOR

Sampling scheme type: Commercial fishing trip

Observation type: SciObsOnShore

Time period of validity: 2022-2027
Short description (max 100 words): sampling scheme aiming at collecting biological samples from commercial landings on-shore about <i>Clupea harengus</i> in Gulf of Riga pelagic trawlers (vessels 12-40m boats). The scheme covers the Gulf of Riga.
Description of the population
<p>Population targeted: PSU is week.</p> <p>Population sampled: On average 2 weeks per month is sampled during the fishing period. No weeks are excluded during the planning.</p> <p>Stratification: There is no stratification.</p>
Sampling design and protocols
<p>Sampling design description: Primary sampling unit (PSU) is time unit of week. Method of PSU selection is NPJS. Sampling scheme follow the Hierarchy 8 in RDBES: sampling time (week) is the primary sampling unit (PSU) and vessel is the secondary sampling units (SSU). Our sampling strategy is that in every month when fishing is ongoing (this depends on the year) subset of week(s) are sampled (PSU=week). Sampling scheme applies to all fractions (crews are interviewed about the presence of whales, while collecting the fish sample).</p> <p>Is the sampling design compliant with the 4S principle?: N</p> <p>Regional coordination: N</p> <p>Link to sampling design documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Compliance with international recommendations: Y</p> <p>Link to sampling protocol documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Compliance with international recommendations: Y</p>
Sampling implementation
<p>Recording of refusal rate: Y</p> <p>Monitoring of sampling progress within the sampling year: On weekly basis online catch and landings reports are used to evaluate the fishing activity and quota exhaustion. This data is compared against the number of samples collected to assure the implementation of the sampling program.</p>
Data capture

Means of data capture: Measuring board, scales, tools to collect otoliths.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: In aggregated form, the data is transmitted to Intercatch and the detailed data to Fishframe or RDB database.

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx>
EMI-UT uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length scatter plots by visual inspection.

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

Sampling design documents in English can be found:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

AtSeaCommercialNEAFC

MS: EST
Region: North Sea and Eastern Arctic, North-East Atlantic
Sampling scheme identifier: AtSeaCommercialNEAFC
Sampling scheme type: Commercial fishing trip
Observation type: SciObsAtSea
Time period of validity: 2022-2027
Short description (max 100 words): sampling scheme aiming at collecting biological samples from commercial landings of selected species at sea by observers. The scheme covers all

Estonian trawlers fishing in North-East Atlantic Fisheries Commission (NEAFC) regulatory area (more precise in ICES areas I and II and XIV).
Description of the population
Population targeted: PSU is fishing trip.
Population sampled: All vessels in the segment are in the draw list.
Stratification: N/A
Sampling design and protocols
Sampling design description: Primary sampling unit (PSU) is fishing trip. Method of PSU selection is SRSWR. Sampling scheme follow the Hierarchy 3 in RDBES.
Is the sampling design compliant with the 4S principle?: Y
Regional coordination: N
Link to sampling design documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Compliance with international recommendations: Y
Link to sampling protocol documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Compliance with international recommendations: Y
Sampling implementation

Recording of refusal rate: Y

Monitoring of sampling progress within the sampling year:

The main concern is recruiting enough observers to be able to cover the sufficient number of long-lasting fishing trips. As the number of vessels is small and companies are interested to have observer onboard there are no big concerns. Changing quotas and short fishing periods could hinder the number of samples collected, thus fishing activity and quota exhaustion is monitored and observers are instructed accordingly on an ongoing basis.

Data capture

Means of data capture: Measuring board, scales, tools to collect otoliths

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database:

EMI-UT

International database: Fishframe/RDB database.

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: Age registering structures are collected but have not been analysed.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight scatter plots by visual inspection.

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

The sampling scheme has been corrected. The PSU is quarter of the year. Fishing trip is the SSU. Method of PSU selection is NPJS. Sampling scheme follow the Hierarchy 3 in RDBES.

Sampling design documents in English can be found:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

AtSeaCommercialNAFO

MS: EST

Region: Other regions (NAFO)

Sampling scheme identifier: AtSeaCommercialNAFO

Sampling scheme type: Commercial fishing trip

Observation type: SciObsAtSea

Time period of validity: 2022-2027

Short description (max 100 words): sampling scheme aiming at collecting biological samples from commercial landings of selected species at sea by observers. The scheme covers all trawling vessels in NAFO areas.

Description of the population

Population targeted: PSU is fishing trip.

Population sampled: Census approach, all vessels fishing in the area are sampled.

Stratification: There is no stratification.

Sampling design and protocols

Sampling design description: Primary sampling unit (PSU) is fishing trip. Method of PSU selection is CENSUS. Sampling scheme follow the Hierarchy 3 in RDBES.

Is the sampling design compliant with the 4S principle?: Y

Regional coordination: N

Link to sampling design documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA (observers on-board are mandatory).

Monitoring of sampling progress within the sampling year: Changing quotas and short fishing periods could hinder the number of samples collected, thus fishing activity and quota exhaustion is monitored and observers are instructed accordingly on an ongoing basis.

Data capture

Means of data capture: Measuring board, scales, tools to collect otoliths.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: Fishframe/RDB database.

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: Age registering structures are collected but have not been analysed.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight scatter plots by visual inspection.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. Sampling design documents in English can be found: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

Parrelectrofishing

MS: EST
Region: Baltic Sea
Sampling scheme identifier: parrelectrofishing
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): sampling scheme aiming at collecting data on <i>Salmo salar</i> and <i>Salmo trutta</i> parr densities in selected rivers using electrofishing. The scheme covers important parr rearing areas. Parr densities of salmonids are sampled annually in 11 fixed rivers inhabited by Atlantic salmon. Parr densities of salmonids are sampled also in 28 to 32 rivers every year which are inhabited mostly by sea trout. 28 rivers are sampled in odd years and 32 in even years.
Description of the population

Population targeted: Target species are Atlantic salmon *Salmo salar* and sea trout *Salmo trutta*. The survey takes place in permanent areas in rivers flowing into the Baltic Sea.

Population sampled: Only salmonid part (*Salmo salar* and *Salmo trutta*) of the river fish assemblage is studied.

Stratification: Parr densities are studied and presented by species (*Salmo salar*, *Salmo trutta*), rivers and areas within the rivers.

Sampling design and protocols

Sampling design description: Monitoring method of studying parr (*Salmo salar* and *Salmo trutta*) densities in rivers is electrofishing. The study sites are fished twice to calculate fishing efficiency and parr densities are presented as individuals per 100 m².

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation:

Method is accredited nationally by the accreditation certificate No L179 (2017) of the Tartu University, Estonian Marine Institute: Assessment of species composition and abundance of salmonids in riverine habitat, KJ I/21; based on Bohlin et al., 1989 and ICES, 2014. Laboratory of the Estonian Marine Institute is accredited against the requirements of standard EVS-EN ISO/IEC 17025:2017 (https://mereinstituut.ut.ee/sites/default/files/mereinstituut/akrediteerimisulatus_2020.pdf).

Compliance with international recommendations: Y

Link to sampling protocol documentation:

Sampling protocol is based on Bohlin *et al* 1989 (Electrofishing – Theory and practice with special emphasis on salmonids. *Hydrobiologia* 173: 9 – 43) article that is modified according to the local needs and this modified methodology is accepted by ICES WGBAST (ICES C.M. 2014/ACOM:08). Sampling design is described in detail in documents of accreditation No L179 (2017) of the Tartu University, Estonian Marine Institute: Assessment of species composition and abundance of salmonids in riverine habitat, KJ I/21.

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: High water levels in river could hinder the sampling. By reserving enough time for sampling, you can find a period when conditions for survey are favourable.

Data capture

Means of data capture: Backpack electrofisher, measuring board, scales.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: NA

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Scales have seldom been collected. If so, then those age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: Estonia takes part in intercalibration workshops.
<https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx>

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight scatter plots by visual inspection.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Sampling design documents in English can be found: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

Smolttrapping

MS: EST

Region: Baltic Sea
Sampling scheme identifier: smolttrapping
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): sampling scheme aiming at collecting data on abundance of descending <i>Salmo salar</i> and <i>Salmo trutta</i> smolts in River Pirita flowing into the Gulf of Finland (SD32).
Description of the population
Population targeted: Target species are Atlantic salmon <i>Salmo salar</i> and sea trout <i>Salmo trutta</i> . The survey takes place in River Pirita, which is designated as index river by ICES.
Population sampled: Only descending salmonid part (<i>Salmo salar</i> and <i>Salmo trutta</i>) of the river fish assemblage is studied.
Stratification: <i>Salmo salar</i> and <i>Salmo trutta</i> smolt abundancy are estimated separately.
Sampling design and protocols
Sampling design description: <i>Salmo salar</i> and <i>Salmo trutta</i> smolt abundancy estimate in river Pirita is done by capture-mark-recapture method. Smolts are caught by trap-net at the river mouth throughout the migration season. Captured smolts are tagged by VIE (visible implant elastomer) and released 2 km upstream from the trap. Some of the tagged smolts will be recaptured during their descent towards the sea. This enables to estimate the overall smolt run size. Only <i>Salmo salar</i> and <i>Salmo trutta</i> smolts are studied, all other fish are released directly.
Is the sampling design compliant with the 4S principle?: NA
Regional coordination: N
Link to sampling design documentation: https://www.agri.ee/kalamajandus-teadusnouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Compliance with international recommendations: Y
Link to sampling protocol documentation: https://www.agri.ee/kalamajandus-teadusnouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi

Compliance with international recommendations: Y
Sampling implementation

Recording of refusal rate: NA
Monitoring of sampling progress within the sampling year: NA
Data capture
Means of data capture: Trap-net, fish tagging equipment, measuring board, scales
Data capture documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Quality checks documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Data storage
National database: EMI-UT
International database: NA
Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Sample storage
Storage description: Scales have seldom been collected. If so, then those age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.
Sample analysis: Estonia takes part in intercalibration workshops. https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx
Data processing
Evaluation of data accuracy (bias and precision): Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Editing and imputation methods: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Quality document associated to a dataset: N
Validation of the final dataset: Deviations are checked by length/weight scatter plots by visual inspection.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Sampling design documents in English can be found:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

Adultcounting

MS: EST
Region: Baltic Sea
Sampling scheme identifier: adultcounting
Sampling scheme type: Diadromous (scientific)
Observation type: EMA water body
Time period of validity: 2022-2027
Short description (max 100 words): sampling scheme aiming at collecting data on abundance of ascending <i>Salmo salar</i> and <i>Salmo trutta</i> adults in River Pirita flowing into the Gulf of Finland (SD32).
Description of the population
<p>Population targeted: Target species are Atlantic salmon <i>Salmo salar</i> and sea trout <i>Salmo trutta</i>. The survey takes place in River Pirita, which is designated as index river by ICES.</p> <p>Population sampled: Ascending Atlantic salmon and sea trout spawners are counted in river Pirita throughout the migration season. River Pirita is designated as index river by ICES. Besides salmon and trout parr density monitoring salmon and sea trout smolt abundancy and spawner abundancy is monitored in river Pirita.</p> <p>Stratification: <i>Salmo salar</i> and <i>Salmo trutta</i> spawners are counted separately.</p>
Sampling design and protocols
<p>Sampling design description: A fish fence (type: resistance board weir) covering the entire width of the river guides fish through an opening that has a fish counter. Only <i>Salmo salar</i> and <i>Salmo trutta</i> adults are counted.</p> <p>Is the sampling design compliant with the 4S principle?: NA</p> <p>Regional coordination: N</p> <p>Link to sampling design documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Compliance with international recommendations: Y</p> <p>Link to sampling protocol documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>

Compliance with international recommendations: Y
Sampling implementation
Recording of refusal rate: NA
Monitoring of sampling progress within the sampling year: NA
Data capture
Means of data capture: Fish counter Vaki Riverwatcher with a camera tunnel.
Data capture documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Quality checks documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Data storage
National database: EMI-UT
International database: NA
Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Sample storage
Storage description: NA
Sample analysis: NA
Data processing
Evaluation of data accuracy (bias and precision): Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Editing and imputation methods: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Quality document associated to a dataset: N
Validation of the final dataset: Videos are analysed by two experts.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. Sampling design documents in English can be found: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi --accordion

CFSeel

MS: EST
Region: Baltic Sea
Sampling scheme identifier: CFSeel
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): sampling scheme aiming at collecting abundance data and biological samples on eel (<i>Anguilla anguilla</i>) in coastal waters of the Baltic Sea. The scheme covers areas SD 28-32.
Description of the population
Population targeted: Target species is <i>Anguilla anguilla</i> and survey takes place in areas SD 28-32. Population sampled: Survey targets yellow eels, silver eels are mostly unreachable for the sampling. Stratification: Survey takes place in 6 permanent coastal areas. Abundance indexes are calculated per area.
Sampling design and protocols

Sampling design description: The fyke nets used are 55 cm high with a semi-circular opening and a leader or wing that is 5 m long. They are made of 17 mm mesh in the arm and 10 mm mesh in the crib of yarn quality no. 210/12 in twisted nylon. Minimum of 80 fyke net days are fished in each of the 6 study areas along the Baltic Sea Estonian coast in July and August. Caught eels are measured for length and weight and otoliths for age reading are collected. Only *Anguilla anguilla*, (mostly yellow eels) are studied, all other fish species are released.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation: Method is accredited nationally by the accreditation certificate No L179 (2017) of the Tartu University, Estonian Marine Institute: Assessment of biodiversity, abundance and biomass of fish assemblages; KJ 1/20; based on EVSEN 14757; Helcom, 2015 (Coastal Fish Guidelines). Laboratory of the Estonian Marine Institute is accredited against the requirements of standard EVS-EN ISO/IEC 17025:2017 (https://mereinstituut.ut.ee/sites/default/files/mereinstituut/akrediteerimisulatus_2020.pdf).

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://helcom.fi/wp-content/uploads/2020/01/HELCOM-Guidelines-for-coastal-fish-monitoring-2019.pdf>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Fyke nets, measuring board, scales.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: NA

Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Sample storage
Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.
Sample analysis: Estonia takes part in intercalibration workshops. https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx
Data processing
Evaluation of data accuracy (bias and precision): Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Editing and imputation methods: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Quality document associated to a dataset: N
Validation of the final dataset: Deviations are checked by length/weight scatter plots by visual inspection.
AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. Sampling design documents in English can be found: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion

Freshwater_eelfykes_scientific

MS: EST
Region: Other
Sampling scheme identifier: freshwater_eelfykes_scientific
Sampling scheme type: Diadromous (scientific)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): Sampling scheme to collect samples from fishery independent (scientific data collection) sources to estimate silver eel escapement from Narva River Basin District.
Description of the population

Population targeted: Target species is *Anguilla anguilla* and survey takes place in Narva River Basin District, most notably Lake Võrtsjärv.

Population sampled: Survey targets yellow eels and silver eels. Males are mostly unreachable due to selectivity of gear.

Stratification: As the eel population is concentrated in Lake Võrtsjärv (85% of restocking takes place in this water body) the samples are collected from only this water body.

Sampling design and protocols

Sampling design description: Fyke nets (mouth opening <3m, mesh size >38mm in the cod end) are used for eel fresh water scientific survey. Survey lasts usually from May until October, depending on water temperature. Fyke nets are controlled in every 3 days. Length (TL=mm), weight (g), age (from otolith), silvering stage (length of the pectoral fins and eye diameter) and infestation with parasites are recorded.

Annual data of eel restocking is collected with average weight (g) and total number of restocked individuals recorded.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation: Sampling design described in Page 87 (Estonian Country Report, Section 4.1):

https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/Fisheries%20Resources%20Steering%20Group/2020/WGEEL_Publications%20with%20multiple%20files/CRs_2020.pdf

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Fyke nets, measuring board, scales, equipment for age reading from otoliths (grinder, microscope)

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: PlutoF <https://plutof.ut.ee>

International database: ICES WGEEL database on http://185.135.126.250:8080/shiny_di/

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Otoliths are stored in Estonian University of Life Sciences. Samples are stored permanently. Access to samples is provided on request. Work is ongoing to insert all information on samples to the database on <https://plutof.ut.ee>

Sample analysis:

<https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2009/WKAREA/WKAREA%202009.pdf>

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://envir.ee/elusloodus-looduskaitse/kalandus/kalanduse-riiklik-andmekogumise-programm-akp#proovivtu-juhendid>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: All datasets are controlled by 3 independent individuals.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan. Sampling design documents in English can be found: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion>

Freshwater_eelfykes_commercial

MS: EST

Region: Other
Sampling scheme identifier: freshwater_eelfykes_commercial
Sampling scheme type: Diadromous (commercial)
Observation type: SciObs water body
Time period of validity: 2022-2027
Short description (max 100 words): Sampling scheme to collect samples from fishery dependent (sampling commercial landings) to estimate silver eel escapement from Narva River Basin District.
Description of the population
<p>Population targeted: PSU is month and SSU is vessel trip, meaning one fishing trip to check all fishing gear. SSU will be selected from a list of trusted fishermen. During a fishing season (April-October), each month one fishing trip is selected for biological data collection, with the main survey area being Lake Võrtsjärv.</p> <p>Population sampled: Population is reachable during the active fyke net fishing period which usually extends from April to October. Both yellow- and silver eels are targeted for sampling in an unsorted catch (meaning the fish selected reflect all specimens the fishermen has caught during that trip).</p> <p>Stratification: As the eel population is concentrated in Lake Võrtsjärv (85% of restocking takes place in this water body) the samples are collected from the same water body.</p>
Sampling design and protocols
<p>Sampling design description: Total length, weight and silvering stage are recorded for the individuals sampled from commercial catches.</p> <p>Is the sampling design compliant with the 4S principle?: NA</p> <p>Regional coordination: N</p> <p>Link to sampling design documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Compliance with international recommendations: N</p> <p>Link to sampling protocol documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Compliance with international recommendations: Y</p>
Sampling implementation

<p>Recording of refusal rate: Y</p> <p>Monitoring of sampling progress within the sampling year: Fishermen are contacted one week prior to following month to make sure of ability to obtain samples. If needed more fishermen will be involved in the program.</p>
<p>Data capture</p>
<p>Means of data capture: Fyke nets, measuring board, scales, equipment for age reading from otoliths (grinder, microscope)</p>
<p>Data capture documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Quality checks documentation: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Data storage</p>
<p>National database: PlutoF https://plutof.ut.ee</p> <p>International database: ICES WGEEL database on http://185.135.126.250:8080/shiny_di/</p> <p>Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Sample storage</p>
<p>Storage description: Otoliths are stored in Estonian University of Life Sciences. Samples are stored permanently. Access to samples is provided on request. Work is ongoing to insert all information on samples to the database on https://plutof.ut.ee</p> <p>Sample analysis: https://www.ices.dk/sites/pub/Publication%20Reports/Expert%20Group%20Report/acom/2009/WKAREA/WKAREA%202009.pdf</p>
<p>Data processing</p>
<p>Evaluation of data accuracy (bias and precision): Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Editing and imputation methods: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Quality document associated to a dataset: N</p> <p>Validation of the final dataset: All datasets are controlled by 3 independent individuals.</p>

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

Sampling design documents in English can be found:

[https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-](https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion)

[kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion](https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi--accordion)

Recreational_catchreports

MS: EST

Region: Baltic Sea

Sampling scheme identifier: recreational_catchreports

Sampling scheme type: Diadromous (recreational)

Observation type: Self water body

Time period of validity: 2022-2027

Short description (max 100 words): Recreational fishermen using commercial gear (gill nets, trap nets, longlines) or fishing for salmon or sea trout in salmonid rivers are all obliged to buy a licence and report back catches. Reports have information about all catches and releases for salmon, sea trout and eel. The scheme covers Estonia's coastal areas (SD 28-32) and salmonid rivers.

Description of the population

Population targeted: All recreational fishermen that fish with gear or areas where having fishing licence is obligatory.

Population sampled: All recreational fishermen that fish with gear or areas where having fishing licence is obligatory.

Stratification: Recreational catches are collected by SD and by rivers.

Sampling design and protocols

Sampling design description: This is a census approach as all recreational fishermen using commercial gear (gill nets, trap nets, longlines) or fishing for salmon or sea trout in salmonid rivers are all obliged to buy a licence and write a report afterwards.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation: NA

Compliance with international recommendations: Y

Link to sampling protocol documentation: NA

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: Y

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: NA

Data capture documentation: NA

Quality checks documentation: NA

Data storage

National database: TEHA (Science and Recreational Fisheries database), overview of yearly catches is available here: <https://envir.ee/elusloodus-looduskaitse/kalandus/harrastuspuuk#statistika>.

International database: NA

Quality checks and data validation documentation: Quality checks are made on database to sort out errors made when entering the data. Fishermen are contacted personally to validate inserted data when inserted information is an outlier (species in uncommon areas, catch volumes etc).

Sample storage

Storage description: NA

Sample analysis: NA

Data processing

Evaluation of data accuracy (bias and precision): NA

Editing and imputation methods: N, imputation is not done.

Quality document associated to a dataset: N

Validation of the final dataset: Quality checks are made on database to sort out errors made when entering the data. Fishermen are contacted personally to validate inserted data when inserted information is an outlier (species in uncommon areas, catch volumes etc).

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

No deviations

Recreational_interview

MS: EST

Region: Baltic Sea

Sampling scheme identifier: Recreational_interview

Sampling scheme type: recreational (off site surveys)

Observation type: EMOnShore

Time period of validity: 2022-2027

Short description (max 100 words): sampling scheme aiming at collecting data about recreational catches and releases for salmon, sea trout, eel and cod by anglers and rod fishermen fishing in the sea. The scheme covers areas 28-32.

Description of the population

Population targeted: The target group of the study is Estonian residents aged 15 and older, who are engaged in recreational angling and rod fishing.

Population sampled: Estonian residents aged 15 and older, younger generation is excluded.

Stratification: Recreational catches are estimated by different coastal areas.

Sampling design and protocols

Sampling design description: Survey was conducted by polling company and used method was CATI (Computer Assisted Telephone Interviewing). The process and the sample size were controlled by the special software (NIPO Fieldwork System for CATI). Filters and rotations of the software are programmed to reduce polling errors. To get better estimates on catch volumes and releases for salmon, sea trout, eel and cod additional interviews were conducted using CAWI (Computer-assisted web interview) method. Outcomes of CAWI interviews were used to have better estimates on catch volumes at sea areas.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation:

<https://www.agri.ee/media/9967/download>

Compliance with international recommendations: Y

Link to sampling protocol documentation:

<https://www.agri.ee/media/9967/download>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: Y

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: CATI (Computer Assisted Telephone Interviewing), CAWI (Computer-assisted web interview) sample size was controlled by the special software (NIPO Fieldwork System for CATI).

Data capture documentation:

<https://www.agri.ee/media/9967/download>

Quality checks documentation:

<https://www.agri.ee/media/9967/download>

Data storage

National database: NA

International database: NA

Quality checks and data validation documentation:

<https://www.agri.ee/media/9967/download>

Sample storage
Storage description: NA
Sample analysis: NA
Data processing
Evaluation of data accuracy (bias and precision): https://www.agri.ee/media/9967/download
Editing and imputation methods: Y, https://www.agri.ee/media/9967/download
Quality document associated to a dataset: N
Validation of the final dataset: The follow-up of the survey was carried out within two days after the end of the survey. The follow-up of the survey was carried out by the project manager of the survey. 10% of the representatives of the target group, with whom the interview was conducted were called again and asked the respondent to comment on various aspects of the interviewer's work. The respondent was asked questions concerning the content of the survey as well as the observance of the survey methodology, adherence to the professional skills. Interviewers whose work was audited were selected on a systematic random basis. Follow-up activities and results were documented.
AR comment: <i>Indicate any deviations. Do not change the text already adopted in the work plan.</i> No deviations. Newest report (incl. description of sampling design) can be found: https://kliimaministerium.ee/sites/default/files/documents/2023-05/2022_HarKal23_aruanne.pdf

BITS_Q4

MS: EST
Region: Baltic Sea
Sampling scheme identifier: BITS_Q4
Sampling scheme type: NA
Observation type: SciObsAtSea
Time period of validity: 2022-2027
Short description (max 100 words): it is the mandatory survey according to Table 1 of the EU MAP Implementing Decision annex. The aim of the survey is monitoring the spatial distribution and abundance of cod and flounder year-classes, and other fish species spatial distribution, taking into consideration the principal hydrological parameters vertical and horizontal variations. The survey is focused on evaluation of the fishing efficiency and abundance of different species and collects materials for the main species biological parameters. The results are used by the ICES WGBFAS for assessment of the cod and flounder stocks.

Description of the population

Population targeted: The main target species are cod and flounder. In Table 1 of the EU MAP Implementing Decision annex target species list of the survey contains the following species: BLL COD DAB FLE HER PLE TUR. But species as BLL, DAB and PLE do not habit or are extremely rare in our survey area and have been caught only on individual occasions or never. EST part of the survey covers SD28.

Population sampled: The main target of the survey is demersal fish assemblage, but pelagic component (Baltic herring) is also analysed.

Stratification: The ICES Subdivisions and depth layers within an ICES Subdivision are used as strata.

Sampling design and protocols

Sampling design description: Total mass and the catch by species are estimated, length composition of main target species, age, sex, individual mass (weight) and maturity samples are collected. Sampling design is described in detail in survey manual:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20\(BITS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20(BITS).pdf)

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: NA

Link to sampling design documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20\(BITS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20(BITS).pdf)

Compliance with international recommendations: Y

Link to sampling protocol documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20\(BITS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20(BITS).pdf)

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Standard trawl type TV-3#520, scales, measuring board, CTD.

Data capture documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20\(BITS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%207%20-%20Manual%20for%20the%20Baltic%20International%20Trawl%20Surveys%20(BITS).pdf)

Quality checks documentation:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-jakalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: DATRAS

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx>
EMI-UT uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: Y, <https://doi.org/10.17895/ices.pub.7580>

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length and other scatter plots by visual inspection.

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

The number of fish and CTD hauls in 2023 was lower than in WP, since actual number and location to be executed by the MS is defined by the ICES WGBIFS.

BIAS

MS: EST
Region: Baltic Sea
Sampling scheme identifier: BIAS
Sampling scheme type: NA
Observation type: SciObsAtSea
Time period of validity: 2022-2027
Short description (max 100 words): it is the mandatory survey according to Table 1 of the EU MAP Implementing Decision annex. The aim of the survey program is to obtain the fisheries-independent information for tuning analytical stock assessment models for Baltic herring and sprat, data are used as indices for assessment purposes.
Description of the population
Population targeted: Target species are Baltic herring and sprat. EST part of the survey covers SD28-32.
Population sampled: Pelagic component of the fish assemblage (Baltic herring and sprat) is studied.
Stratification: The stratification is based on ICES statistical rectangles.
Sampling design and protocols
Sampling design description: The acoustic surveys cover the total area of ICES Division III and site allocation is done by the Baltic International Fish Survey Working Group (WGBIFS), thus each country has a mandatory responsible area. The area is limited inshore by the 10 m depth line. The standard equipment used for the survey is the Simrad EK/EY-60 echosounder and the standard frequency is 38 kHz. Baltic International Acoustic Survey (BIAS) is carried out in September/October. Sampling design is described in detail in survey manual:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SI SP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: NA

Link to sampling design documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SI SP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Compliance with international recommendations: Y

Link to sampling protocol documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SI SP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Standard trawl, Simrad EK/EY-60 echosounder, scales, measuring board, CTD.

Data capture documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SI SP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Quality checks documentation:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: ICES Acoustic trawl Surveys Database

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus->

[nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi](https://www.agri.ee/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi)

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx>
EMI-UT uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: Y, <https://doi.org/10.17895/ices.pub.7581>

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length and other scatter plots by visual inspection.

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

No deviations.

GRAHS

MS: EST

Region: Baltic Sea

Sampling scheme identifier: GRAHS

Sampling scheme type: NA

Observation type: SciObsAtSea

Time period of validity: 2022-2027

Short description (max 100 words): it is the mandatory survey according to Table 1 of the EU MAP Implementing Decision annex. The aim of the survey program is to obtain the fisheries-independent information for tuning analytical stock assessment model for Baltic herring in the Gulf of Riga, data is used as indices for assessment purposes. Survey takes place in the Gulf of Riga.

Description of the population

Population targeted: Target species is Baltic herring. Survey covers the whole Gulf of Riga

Population sampled: Pelagic fish species Baltic herring is studied.

Stratification: NA

Sampling design and protocols

Sampling design description: The acoustic surveys cover the total area of the Gulf of Riga. The survey is carried out jointly by the Latvian and Estonian scientists on the chartered Latvian fishing vessel. The standard equipment used for the survey is the Simrad EK/EY-60 echosounder and the standard frequency is 38 kHz. GRAHS is carried out in July-August annually in order to cover the period after main spawning season when most of the stock has left the near-coast spawning grounds. Sampling design is described in detail in survey manual:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: NA

Link to sampling design documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Compliance with international recommendations: Y

Link to sampling protocol documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Standard trawl, Simrad EK/EY-60 echosounder, scales, measuring board, CTD

Data capture documentation:

[https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20\(SISP\)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20\(IBAS\).pdf](https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SISP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf)

Quality checks documentation:

<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: ICES Acoustic trawl Surveys Database

Quality checks and data validation documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx>
EMI-UT uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: Y, <https://doi.org/10.17895/ices.pub.7581>

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length and other scatter plots by visual inspection.

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

Survey duration was shorter due to good weather conditions. Since the fish aggregations were concentrated in the northern part of gulf and some planned trawling areas had no fish indications, the number of executed hauls was reduced to shorten fuel costs.

SPRAS

MS: EST
Region: Baltic Sea
Sampling scheme identifier: SPRAS
Sampling scheme type: NA
Observation type: SciObsAtSea
Time period of validity: 2022-2027
Short description (max 100 words): it is the mandatory survey according to Table 1 of the EU MAP Implementing Decision annex. The aim of the survey program is to obtain the fisheries-independent information for tuning analytical stock assessment models for sprat, data are used as indices for assessment purposes.
Description of the population
Population targeted: Target species is sprat (<i>Sprattus spattus</i>). EST part of the survey covers SD28-32.
Population sampled: Pelagic fish species sprat (<i>Sprattus spattus</i>) is studied.
Stratification: The stratification is based on ICES statistical rectangles.
Sampling design and protocols
Sampling design description: The acoustic surveys cover the total area of ICES Division III and area allocations is done by the Baltic International Fish Survey Working Group (WGBIFS), thus each country has a mandatory responsible area. The area is limited inshore by the 10 m depth line. The standard equipment used for the survey is the Simrad EK/EY-60 echosounder and the standard frequency is 38 kHz. Baltic International Acoustic Survey (SPRAS) is carried out in May. Sampling design is described in detail in survey manual: https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf
Is the sampling design compliant with the 4S principle?: NA
Regional coordination: NA
Link to sampling design documentation: https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf
Compliance with international recommendations: Y
Link to sampling protocol documentation: https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf

<p>SP/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf</p> <p>Compliance with international recommendations: N</p>
<p>Sampling implementation</p>
<p>Recording of refusal rate: NA</p>
<p>Monitoring of sampling progress within the sampling year: NA</p>
<p>Data capture</p>
<p>Means of data capture: Standard trawl, Simrad EK/EY-60 echosounder, scales, measuring board, CTD.</p> <p>Data capture documentation: https://www.ices.dk/sites/pub/Publication%20Reports/ICES%20Survey%20Protocols%20(SI%20SP)/SISP%208%20-%20Manual%20of%20International%20Baltic%20Acoustic%20Surveys%20(IBAS).pdf</p>
<p>Quality checks documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Data storage</p>
<p>National database: EMI-UT</p> <p>International database: ICES Acoustic trawl Surveys Database</p> <p>Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p>
<p>Sample storage</p>
<p>Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.</p> <p>Sample analysis: https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx EMI-UT uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.</p>
<p>Data processing</p>

Evaluation of data accuracy (bias and precision): Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y, <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: Y, <https://doi.org/10.17895/ices.pub.7580>

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length and other scatter plots by visual inspection

AR comment: *Indicate any deviations. Do not change the text already adopted in the work plan.*

No deviations.

EFLS

MS: EST

Region: Baltic Sea

Sampling scheme identifier: EFLS

Sampling scheme type: NA

Observation type: SciObsAtSea

Time period of validity: 2022-2027

Short description (max 100 words): national survey has been conducted annually since 1947 to study the distribution and abundance of fish larvae and juveniles with the aim to provide primary information on herring yearclass abundance and to understand the affecting environmental background.

Description of the population

Population targeted: The main target species is spring spawning Baltic herring. The survey covers NE part of the Gulf of Riga.

Population sampled: Ichthyoplankton and mezozooplankton communities. Larvae and juveniles of all fish species are identified, measured and counted in NE part of Gulf of Riga.

Stratification: Fixed stations in NE part of Gulf of Riga.

Sampling design and protocols

Sampling design description: Nine fixed stations are visited weekly (May-July). Hensen larval fish net is used for 10 min. hauls in commercially important fish spawning and nursery grounds. Larvae are identified, measured and counted. Plankton samples and environmental data are collected and analysed.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation: Details of the survey design is described in following paper: <https://www.sciencedirect.com/science/article/abs/pii/S016578361000281X>
Mesozooplankton samples are collected according to HELCOM manual:
<https://helcom.fi/media/publications/Guidelines-for-monitoring-of-mesozooplankton.pdf>

Compliance with international recommendations: Y. Details of the survey design is described in following paper:
<https://www.sciencedirect.com/science/article/abs/pii/S016578361000281X>
Mesozooplankton samples are collected according to HELCOM manual:
<https://helcom.fi/media/publications/Guidelines-for-monitoring-of-mesozooplankton.pdf>

Link to sampling protocol documentation: Sampling is described in detail in paper:
<https://www.sciencedirect.com/science/article/abs/pii/S016578361000281X>
Mesozooplankton samples are collected according to HELCOM manual:
<https://helcom.fi/media/publications/Guidelines-for-monitoring-of-mesozooplankton.pdf>

Compliance with international recommendations: Y, Sampling is described in detail in paper: <https://www.sciencedirect.com/science/article/abs/pii/S016578361000281X>
Mesozooplankton samples are collected according to HELCOM manual:
<https://helcom.fi/media/publications/Guidelines-for-monitoring-of-mesozooplankton.pdf>

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Binocular Stemi 2000-CS, CTD, Hensen net, mesozooplankton WP-2 net equipped with flowmeters

Data capture documentation: <https://www.agri.ee/media/11646/download>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage
National database: EMI-UT
International database: ICES WGALES https://eggsandlarvae.ices.dk/Download.aspx
Quality checks and data validation documentation: https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Sample storage
Storage description: Fish larvae and mezozooplankton samples are stored in 4% formaldehyde solution. Samples are deposited in Estonian Marine Institute storage room for 5 yr. Responsible person takes care of a maintenance, conservation and accessibility to samples.
Sample analysis: https://www.agri.ee/media/11646/download https://www.sciencedirect.com/science/article/abs/pii/S016578361000281X https://helcom.fi/media/publications/Guidelines-for-monitoring-of-mesozooplankton.pdf Kazanova, I. 1953. Apredelitel ikrõ i litsinok rõb Baltiskovo morja i jevo salivov. Trudõ Vsjesojusnovo naustno-isledovatelskova instituta marskova rõbnava hasjastva I okenografi (VNIRO)
Data processing
Evaluation of data accuracy (bias and precision): https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Editing and imputation methods: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi
Quality document associated to a dataset: N
Validation of the final dataset: The end-user of datasets is Fisheries department of Estonian Ministry of Environment that applies collected dataset implementing spatio-temporal fishing regulations and marine management. Validation and quality checks are performed in regular manner during annual reporting phase by checking all submitted datasets step-by-step interactively with researchers.
AR comment: <i>Indicate any deviations. Do not change the text already adopted in the work plan.</i> Planned days at sea and number of planned sampling activities are the maximum numbers, in consideration of trip per week from May to August. The actual trip number depends on the hatching and development of larvae. Surveys end when herring larvae reach the developmental stage at which they can avoid the gear. It happens now earlier because of increased water temperatures during spawning and hatching season which results in a faster growth rate and as high water temperatures are lethal to yolk-sac stage larvae then herring hatched in late June or early July do not survive. So since mid-July there are no herring larvae to sample. We have not detected any change in the start of herring spawning.

GORFS

MS: EST
Region: Baltic Sea
Sampling scheme identifier: GORFS
Sampling scheme type: NA
Observation type: SciObsAtSea
Time period of validity: 2022-2027
Short description (max 100 words): Trawl survey in the Pärnu Bay was conducted already in earlier decades (since 1950s) and resumed in 2009. It is a national survey aiming at collecting of fisheries independent data for tuning the pikeperch and perch VPA assessment and collecting information about abundance and biological parameters of other fish species, including commercially important species like smelt and cyprinids.
Description of the population
<p>Population targeted: The main target species are perch (<i>Perca fluviatilis</i>) and pikeperch (<i>Sander lucioperca</i>). Six fixed trawl transects are situated three to eight km from shore (water depth five to nine m) to cover the entire length of the Pärnu Bay.</p> <p>Population sampled: In shallow waters of the Pärnu bay demersal and pelagic part of the fish community are involved. Data on the abundance and biological parameters of fish species in the temporal fish assemblages are collected.</p> <p>Stratification: Fixed stations in Pärnu bay.</p>
Sampling design and protocols

Sampling design description: Research vessel of the Estonian Marine Institute AURELIE is used for trawling. The trawl (working depth 0.3 m from the bottom) is pulled with the speed of 3 knots for 30 minutes. The trawl mouth is 2 m high and 6 m wide, distance between doors is 20 m and maximum distance between the 8.2 m long trawl wings is 12 m. Mesh size is 60 mm (knot to knot), at the tip of the trawl wings, 45 mm at the trawl mouth and decreases gradually to 10 mm at the codend. All 6 fixed stations are fished in May, September, October, November and December.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation: <https://helcom.fi/wp-content/uploads/2020/01/HELCOM-Guidelines-for-coastal-fish-monitoring-2019.pdf>

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://helcom.fi/wp-content/uploads/2020/01/HELCOM-Guidelines-for-coastal-fish-monitoring-2019.pdf>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Standard trawl, Simrad PI50 trawl monitoring system, scales, measuring board.

Data capture documentation: <https://helcom.fi/wp-content/uploads/2020/01/HELCOM-Guidelines-for-coastal-fish-monitoring-2019.pdf>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: NA

Quality checks and data validation documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage
<p>Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.</p> <p>Sample analysis: Thoresson, G. 1996. Guidelines for coastal monitoring. Kustrapport 1: 1-35. https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx Estonia uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops</p>
Data processing
<p>Evaluation of data accuracy (bias and precision): Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Editing and imputation methods: Y, https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi</p> <p>Quality document associated to a dataset: N</p> <p>Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length scatter plots by visual inspection.</p> <p>AR comment: <i>Indicate any deviations. Do not change the text already adopted in the work plan.</i></p> <p>No deviations.</p>

CFS

MS: EST
Region: Baltic Sea
Sampling scheme identifier: CFS
Sampling scheme type: NA
Observation type: SciObsAtSea
Time period of validity: 2022-2027

Short description (max 100 words): National annual survey using standardised gill net series started in 1992, now in 10 fixed areas. Collected information (CPUE, age and length distribution, etc) form the basis for advice for commercially important stocks (perch, pikeperch, flounder, cod, whitefish, cyprinids) including fisheries independent data for tuning the flounder, pikeperch and perch analytical assessment.

Description of the population

Population targeted: The main target species are perch, flounder, pikeperch and cod. The survey takes place in 10 fixed areas in SD28, 29 and 32.

Population sampled: Demersal part of the fish community is targeted but pelagic fish community is also partly involved. All caught fish species are identified, measured and weighted, also those not namely listed in the National WP or in Table 1 of the EU MAP Delegated Decision annex.

Stratification: S28, SD29 and SD32 overall 10 fixed areas.

Sampling design and protocols

Sampling design description: Annual gill-net survey in defined areas along the Estonian coast, in fixed (Hiiumaa, Kõiguste and Küdema) or random stations. Each station consists of a series of gill nets of fixed mesh sizes and construction. 6-72 stations are fished in each area.

Is the sampling design compliant with the 4S principle?: NA

Regional coordination: N

Link to sampling design documentation: Method is accredited nationally by the accreditation certificate No L179 (2017) of the Tartu University, Estonian Marine Institute: Assessment of biodiversity, abundance and biomass of fish assemblages; KJ 1/20; based on EVSEN 14757; Helcom, 2015 (Coastal Fish Guidelines). Laboratory of the Estonian Marine Institute is accredited against the requirements of standard EVS-EN ISO/IEC 17025:2017 (https://mereinstituut.ut.ee/sites/default/files/mereinstituut/akrediteerimisulatus_2020.pdf).

Compliance with international recommendations: Y

Link to sampling protocol documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Compliance with international recommendations: Y

Sampling implementation

Recording of refusal rate: NA

Monitoring of sampling progress within the sampling year: NA

Data capture

Means of data capture: Standardized gill nets, measuring board, scales.

Data capture documentation: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality checks documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Data storage

National database: EMI-UT

International database: Data is partly in HELCOM Coastal fish core indicator database (COOL), [http://bio.helcom.fi/apex/f?p=108:5:::~::](http://bio.helcom.fi/apex/f?p=108:5:::)

Quality checks and data validation documentation:
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Sample storage

Storage description: Age registering samples are stored in EMI-UT archive permanently. Archive is accessible to the staff of EMI-UT and to another scientist on request. The work is ongoing to digitize the data in archive and make the information about the stored materials available through the Web.

Sample analysis: Thoresson, G. 1996. Guidelines for coastal monitoring. Kustrapport 1: 1-35. <https://www.ices.dk/community/Pages/PGCCDBS-doc-repository.aspx> Estonia uses SmartDots age reading platform, which facilitates age readings based on otolith images and takes part in intercalibration workshops.

Data processing

Evaluation of data accuracy (bias and precision): Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Editing and imputation methods: Y,
<https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Quality document associated to a dataset: N

Validation of the final dataset: Deviations are checked by length/weight, age/weight, age/length scatter plots by visual inspection.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

No deviations.

ANNEX 1.2 - QUALITY REPORT FOR SOCIOECONOMIC DATA SAMPLING SCHEME

The quality report fulfils Article 6 (3) (d) of Regulation (EU) 2017/1004. This document is intended to specify data to be collected under Chapter II, points 3, 5, 6, and 7 of the Delegated Decision annex: 'Socioeconomic data on fisheries, aquaculture and any complementary data collection of fishing activity and fish processing'. Use this annex to describe quality aspects of the data collection process (design, sampling implementation, data capture, data storage and data processing etc.). The annex should be filled for each sampling scheme. Where applicable, use the handbook on sampling design (Deliverable 2.1 from MARE/2016/22 SECFISH study), available on the DCF website.

SocioEconomic Fisheries data: Census

<p>Survey Specifications</p> <p><i>'Sector name' refers to socio economic data on fisheries, aquaculture and any complementary data collection for fishing activities and processing, as in the EU MAP Delegated Decision annex.</i></p> <p><i>'Sampling scheme' refers to the survey technique: by census, by sampling, random or non-random, other (with explanation). If sampling, then outline sampling design.</i></p> <p><i>'Variables' refer to Tables 7, 9 and 10 of the EU MAP Delegated Decision annex.</i></p> <p><i>'Supra region' refers to Table 2 of the EU MAP Implementing Decision annex. If the sampling scheme is the same in all supra regions, put 'All supra regions'.</i></p>
<p>Sector name(s): Fisheries data</p>
<p>Sampling scheme: census</p>
<p>Variables: Table 7 and 9 of the EU MAP Delegated Decision annex</p>
<p>Supra region(s): Baltic Sea; North Sea; Eastern Arctic; NAFO; extended North Western waters (ICES areas 5, 6 and 7) and extended South Western waters (ICES areas 10, 12 and 14)</p>
<p>Survey planning</p>
<p>Sampling scheme applies to all trawlers and those coastal boats that catch at least 100t per year (based on previous year's catches)</p>
<p>Survey design and strategy</p> <p>Describe how the sample sizes were determined: Complete enumeration is applied.</p> <p>Describe survey methods and distribution, e.g. questionnaire forms by post, by email, on website, by phone, access to other datasets, etc.</p> <p>Data are collected and the submission of questionnaires is monitored through eSTAT (the web channel for electronic data submission). The questionnaires have been designed for independent completion in eSTAT and include instructions and controls. The questionnaires and information about data submission are available on the website of Statistics Estonia in the section Questionnaires.</p> <p>Describe the role of auxiliary information, if any, in the strategy, e.g. for validation, cross referencing, as a fall-back data source, etc.</p>

We use fishing data, annual statistics and tax office data for additional information.

Estimation design

Describe method of calculating population estimate from sample.
Describe method of calculating derived data, e.g. imputed values.
Describe treatment of non-responses.

The response rate has been 100%, so we do not have problem with non-response. System is that we send the reminders 5 days before the deadline, and 3, 7 and 21 days after the deadline. In case of non-submission penalty system may be used.

Error checks

Describe potential errors and how and where in the process these are detected, avoided or eliminated, e.g. data duplication, double counting, respondent error, upload error, processing error, etc.

The input data are checked and, if necessary, corrected according to specified rules, and the compliance of calculated statistics with the quality requirements is checked as well: arithmetic and qualitative controls, comparisons are made with previous periods, internal coherence, the coverage of the population and the response rate are checked.

Data storage and documentation

Describe how the data are stored.
Provide a link to the webpage where additional methodological documentation can be found, if any.

The data are in the official database of Statistical Estonia, but are not publicly available. Methodological documentation is available: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Revision

Describe the frequency of the methodology review e.g. revision of segmentation, survey method per segment, per variable, etc.

There are no planned revisions. The data may be revised if the methodology is modified, errors are discovered, new or better data become available. The data revision policy and notification of corrections are described in the section Principles of dissemination of official statistics of the website of Statistics Estonia.

Confidentiality

Are procedures for confidential data handling in place and documented?
Are protocols to enforce confidentiality between DCF partners in place and documented?
Are protocols to enforce confidentiality with external users in place and documented?
Are there any issues with publication of data due to confidentiality reasons? Provide an explanation.

The dissemination of data collected for the purpose of producing official statistics is guided by the requirements provided for in § 32, § 34, § 35, § 38 of the Official Statistics Act. The

treatment of confidential data is regulated by the Procedure for Protection of Data Collected and Processed by Statistics Estonia (in Estonian). See more details on the website of Statistics Estonia in the section Õigusaktid.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

No deviations from the work plan.

SocioEconomic Fisheries data: PSS

Survey Specifications

'Sector name' refers to socio economic data on fisheries, aquaculture and any complementary data collection for fishing activities and processing, as in the EU MAP Delegated Decision annex.

'Sampling scheme' refers to the survey technique: by census, by sampling, random or non-random, other (with explanation). If sampling, then outline sampling design.

'Variables' refer to Tables 7, 9 and 10 of the EU MAP Delegated Decision annex.

'Supra region' refers to Table 2 of the EU MAP Implementing Decision annex. If the sampling scheme is the same in all supra regions, put 'All supra regions'.

Sector name(s): Fisheries data

Sampling scheme: Stratified simple random sample

Variables: Table 7 and 9 of the EU MAP Delegated Decision annex

Supra region(s): Baltic Sea; North Sea; Eastern Arctic; NAFO; extended North Western waters (ICES areas 5, 6 and 7) and extended South Western waters (ICES areas 10, 12 and 14)

Survey planning

Sampling scheme applies to those coastal boats that catch less than 100t per year (based on previous year's catches). The population is divided into two strata: 1) owners of coastal boats who caught less than 10 tons (stratum 1) and 2) owners of coastal boats who caught from 10 up to 100 tons (stratum 2) in previous year. A simple random sample is drawn from the both stratum and different regions are taken into account in the final sample.

Survey design and strategy

List data sources, e.g. interviews, registers, log books, sales notes, VMS, financial accounts, etc.
 For determining the sample fleet register, log books and sales note information is used.
 Information from entities that end up in a sample is collected via online tool described below.

Describe how the sample sizes were determined.

In these two strata described above sample sizes were chosen as follows:

- Stratum 1 (boat owners with catch less than 10t in previous year): 8% of units are selected
- Stratum 2 (boat owners with catch between 10t to 100t in previous year): 50% of units are selected.

Describe survey methods and distribution, e.g. questionnaire forms by post, by email, on website, by phone, access to other datasets, etc.

Data are collected and the submission of questionnaires is monitored through eSTAT (the web channel for electronic data submission). The questionnaires have been designed for independent completion in eSTAT and include instructions and controls. The questionnaires and information about data submission are available on the website of Statistics Estonia in the section Questionnaires.

Describe the role of auxiliary information, if any, in the strategy, e.g. for validation, cross referencing, as a fall-back data source, etc.

We use fishing data, annual statistics and tax office data for additional information.

Estimation design

Describe method of calculating population estimate from sample.

For calculating an estimate \hat{t}_y of a population total t_y of a variable y the Horvitz-Thompson

estimator is used, i.e. $\hat{t}_y = \sum_{h=1}^H \frac{N_H}{n_h} \sum_{i_h=1}^{n_h} y_{i_h}$, where H is the number of strata, N_h is the population size of the stratum h , n_h is the sample size of the stratum h and $\sum_{i_h=1}^{n_h} y_{i_h}$ is the

sample total of the variable y . Information about variables (all, except variable group Fleet, Effort and Number of fishing enterprises/units) are collected from fishing enterprises.

Describe treatment of non-responses. The response rate has been almost 100%, so we do not have problem with non-response. System is that Statistics Estonia sends the remainders 5 days before the deadline, and 3, 7 and 21 days after the deadline. In case of non-submission penalty system may be used. In total imputation is not applied, effective sample size is just smaller if any entity fails to respond.

Error checks

Describe potential errors and how and where in the process these are detected, avoided or eliminated, e.g. data duplication, double counting, respondent error, upload error, processing error, etc.

The input data are checked and, if necessary, corrected according to specified rules, and the compliance of calculated statistics with the quality requirements is checked as well: arithmetic and qualitative controls, comparisons are made with previous periods, internal coherence, the coverage of the population and the response rate are checked.

Data storage and documentation

Describe how the data are stored.

Provide a link to the webpage where additional methodological documentation can be found, if any.

The data are in the official database of Statistical Estonia, but are not publicly available.

Methodological documentation is available: <https://www.agri.ee/kalamajandus-teadus-nouanne/kalamajandus-ja-kalapuuk/riiklik-andmekogumisprogramm#sampling-methodologi>

Revision

Describe the frequency of the methodology review e.g. revision of segmentation, survey method per segment, per variable, etc.

There are no planned revisions. The data may be revised if the methodology is modified, errors are discovered, new or better data become available. The data revision policy and notification of corrections are described in the section Principles of dissemination of official statistics of the website of Statistics Estonia.

Confidentiality

Are procedures for confidential data handling in place and documented?

Are protocols to enforce confidentiality between DCF partners in place and documented?

Are protocols to enforce confidentiality with external users in place and documented?

Are there any issues with publication of data due to confidentiality reasons? Provide an explanation.

The dissemination of data collected for the purpose of producing official statistics is guided by the requirements provided for in § 32, § 34, § 35, § 38 of the Official Statistics Act. The treatment of confidential data is regulated by the Procedure for Protection of Data Collected and Processed by Statistics Estonia (in Estonian). See more details on the website of Statistics Estonia in the section Õigusaktid.

AR comment: Indicate any deviations. Do not change the text already adopted in the work plan.

No deviations from the work plan.